



The  
Patent  
Office

PCT/US 98/10951

ROUS 28 SEP 1998

The Patent Office  
Cardiff Road  
Newport  
Gwent  
NP9 1RH

REC'D	20 OCT 1998
WIPO	PCT

UK 9800680.2

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation and Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words public limited company may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Signed

*Answee*

Dated

23 September 1998







**The  
Patent  
Office**

**1/77**

**Request for grant of a patent**

(See notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

**The Patent Office**

**Cardiff Road  
Newport  
Gwent NP9 1RH**

1. Your reference

19964PV2

15JAN98 E330401-1 D02639  
P01/7700 25.00 - 9800680.2

2. Patent application number

(The Patent Office will fill in this part)

**9800680.2**

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Merck & Co., Inc.  
P. O. Box 2000  
Rahway, New Jersey 07065-0900  
U.S.A.

Patents ADP number (if you know it)

00597674001

If the applicant is a corporate body, give the country/state of its incorporation

New Jersey, USA

4. Title of the invention

Biarylalkanoic acids as cell adhesion inhibitors

5. Name of your agent (if you have one)

Mr. I. J. Hiscock

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Merck & Co., Inc.  
European Patent Department  
Terlings Park  
Eastwick Road  
Harlow  
Essex CM20 2QR

Patents ADP number (if you know it)

06546683001 ✓

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority Application number  
(if you know it)

Date of filing  
(day/month/year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(day/month/year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

Yes

- a) any applicant named in part 3 is not an inventor, or
  - b) there is an inventor who is not named as an applicant, or
  - c) any named applicant is a corporate body.
- See note (d))

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document


Continuation sheets of this form	0
Description	61
Claim(s)	12
Abstract	1
Drawing(s)	0



10. If you are also filing any of the following, state how many against each item.

Priority documents	-
Translations of priority documents	-
Statement of inventorship and right to grant of a patent (Patents Form 7/77)	-
Request for preliminary examination and search (Patents Form 9/77)	-
Request for substantive examination (Patents Form 10/77)	-
Any other documents (please specify)	Fee Sheet

11. I/We request the grant of a patent on the basis of this application.

Signature  Date 12 January 1998  
Mr. I. J. Hiscock

12. Name and daytime telephone number of person to contact in the United Kingdom
- |                   |              |
|-------------------|--------------|
| Mr. I. J. Hiscock | 01279 440175 |
|-------------------|--------------|

**Warning**

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

**Notes**

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

## TITLE OF THE INVENTION

## BIARYLALKANOIC ACIDS AS CELL ADHESION INHIBITORS

### BACKGROUND OF THE INVENTION

5           The present invention relates to novel substituted biarylalkanoic acid derivatives which are useful for the inhibition and prevention of leukocyte adhesion and leukocyte adhesion-mediated pathologies. This invention also relates to compositions containing such compounds and methods of treatment using such compounds.

10           Many physiological processes require that cells come into close contact with other cells and/or extracellular matrix. Such adhesion events may be required for cell activation, migration, proliferation and differentiation. Cell-cell and cell-matrix interactions are mediated through several families of cell adhesion molecules (CAMs) including  
15 the selectins, integrins, cadherins and immunoglobulins. CAMs play an essential role in both normal and pathophysiological processes. Therefore, the targetting of specific and relevant CAMs in certain disease conditions without interfering with normal cellular functions is essential for an effective and safe therapeutic agent that inhibits cell-cell  
20 and cell-matrix interactions.

          The integrin superfamily is made up of structurally and functionally related glycoproteins consisting of  $\alpha$  and  $\beta$  heterodimeric, transmembrane receptor molecules found in various combinations on nearly every mammalian cell type (for reviews see: E. C. Butcher, Cell,  
25 67, 1033 (1991); T. A. Springer, Cell, 76, 301 (1994); D. Cox et al., "The Pharmacology of the Integrins." Medicinal Research Rev. 14, 195 (1994) and V. W. Engleman et al., "Cell Adhesion Integrins as Pharmaceutical Targets." in Ann. Repts. in Medicinal Chemistry, Vol. 31, J. A. Bristol, Ed.; Acad. Press, NY, 1996, p. 191).

30           VLA-4 ("very late antigen-4"; CD49d/CD29; or  $\alpha 4\beta 1$ ) is an integrin expressed on all leukocytes, except platelets and mature neutrophils, and is a key mediator of the cell-cell and cell-matrix interactions of leukocytes (see M. E. Hemler, "VLA Proteins in the Integrin Family: Structures, Functions, and Their Role on Leukocytes."

Ann. Rev. Immunol. 8, 365 (1990)). The ligands for VLA-4 include vascular cell adhesion molecule-1 (VCAM-1) and the CS-1 domain of fibronectin (FN). VCAM-1 is a member of the Ig superfamily and is expressed *in vivo* on endothelial cells at sites of inflammation and on dendritic and macrophage-like cells. (See R. Lobb et al. "Vascular Cell Adhesion Molecule 1." in Cellular and Molecular Mechanisms of Inflammation, C. G. Cochrane and M. A. Gimbrone, Eds.; Acad. Press, San Diego, 1993, p. 151.) VCAM-1 is produced by vascular endothelial cells in response to pro-inflammatory cytokines (See A. J. H. Gearing and W. Newman, "Circulating adhesion molecules in disease.", Immunol. Today, 14, 506 (1993). The CS-1 domain is a 25 amino acid sequence that arises by alternative splicing within a region of fibronectin. (For a review, see R. O. Hynes "Fibronectins.", Springer-Verlag, NY, 1990.) A role for VLA-4/CS-1 interactions in inflammatory conditions has been proposed (see M. J. Elices, "The integrin  $\alpha_4\beta_1$  (VLA-4) as a therapeutic target" in Cell Adhesion and Human Disease, Ciba Found. Symp., John Wiley & Sons, NY, 1995, p. 79).

$\alpha_4\beta_7$  (also referred to as LPAM-1 and  $\alpha_4\beta_p$ ) is an integrin expressed on leukocytes and is a key mediator of leukocyte trafficking and homing in the gastrointestinal tract (see C. M. Parker et al., Proc. Natl. Acad. Sci. USA, 89, 1924 (1992)). The ligands for  $\alpha_4\beta_7$  include mucosal addressing cell adhesion molecule-1 (MadCAM-1) and, upon activation of  $\alpha_4\beta_7$ , VCAM-1 and fibronectin (Fn). MadCAM-1 is a member of the Ig superfamily and is expressed *in vivo* on endothelial cells of gut-associated mucosal tissues of the small and large intestine ("Peyer's Patches") and lactating mammary glands. (See M. J. Briskin et al., Nature, 363, 461 (1993); A. Hamann et al., J. Immunol., 152, 3282 (1994)). MadCAM-1 can be induced *in vitro* by proinflammatory stimuli (See E. E. Sikorski et al. J. Immunol., 151, 5239 (1993)). MadCAM-1 is selectively expressed at sites of lymphocyte extravasation and specifically binds to the integrin,  $\alpha_4\beta_7$ .

Neutralizing anti- $\alpha_4$  antibodies or blocking peptides that inhibit the interaction between VLA-4 and/or  $\alpha_4\beta_7$  and their ligands

have proven efficacious both prophylactically and therapeutically in several animal models of disease, including i) experimental allergic encephalomyelitis, a model of neuronal demyelination resembling multiple sclerosis (for example, see T. Yednock et al., "Prevention of experimental autoimmune encephalomyelitis by antibodies against  $\alpha_4\beta_1$  integrin." Nature, 356, 63 (1993) and E. Keszthelyi et al., "Evidence for a prolonged role of  $\alpha_4$  integrin throughout active experimental allergic encephalomyelitis." Neurology, 47, 1053 (1996)); ii) bronchial hyperresponsiveness in sheep and guinea pigs as models for the various phases of asthma (for example, see W. M. Abraham et al., " $\alpha_4$ -Integrins mediate antigen-induced late bronchial responses and prolonged airway hyperresponsiveness in sheep." J. Clin. Invest. 93, 776 (1993) and A. A. Y. Milne and P. P. Piper, "Role of VLA-4 integrin in leucocyte recruitment and bronchial hyperresponsiveness in the guinea-pig." Eur. J. Pharmacol., 282, 243 (1995)); iii) adjuvant-induced arthritis in rats as a model of inflammatory arthritis (see C. Barbadillo et al., "Anti-VLA-4 mAb prevents adjuvant arthritis in Lewis rats." Arthr. Rheuma. (Suppl.), 36 95 (1993) and D. Seiffge, "Protective effects of monoclonal antibody to VLA-4 on leukocyte adhesion and course of disease in adjuvant arthritis in rats." J. Rheumatol., 23, 12 (1996)); iv) adoptive autoimmune diabetes in the NOD mouse (see J. L. Baron et al., "The pathogenesis of adoptive murine autoimmune diabetes requires an interaction between  $\alpha_4$ -integrins and vascular cell adhesion molecule-1." J. Clin. Invest., 93, 1700 (1994), A. Jakubowski et al., "Vascular cell adhesion molecule-Ig fusion protein selectively targets activated  $\alpha_4$ -integrin receptors in vivo: Inhibition of autoimmune diabetes in an adoptive transfer model in nonobese diabetic mice." J. Immunol., 155, 938 (1995), and X. D. Yang et al., "Involvement of beta 7 integrin and mucosal addressin cell adhesion molecule-1 (MadCAM-1) in the development of diabetes in nonobese diabetic mice", Diabetes, 46, 1542 (1997)); v) cardiac allograft survival in mice as a model of organ transplantation (see M. Isobe et al., "Effect of anti-VCAM-1 and anti-VLA-4 monoclonal antibodies on cardiac allograft survival and response to soluble antigens in mice." Transplant. Proc., 26, 867 (1994) and S.

- Molossi et al., "Blockade of very late antigen-4 integrin binding to fibronectin with connecting segment-1 peptide reduces accelerated coronary arteriopathy in rabbit cardiac allografts." J. Clin. Invest., **95**, 2601 (1995)); vi) spontaneous chronic colitis in cotton-top tamarins which resembles human ulcerative colitis, a form of inflammatory bowel disease (see D. K. Podolsky et al., "Attenuation of colitis in the Cotton-top tamarin by anti- $\alpha_4$  integrin monoclonal antibody.", J. Clin. Invest., **92**, 372 (1993)); vii) contact hypersensitivity models as a model for skin allergic reactions (see T. A. Ferguson and T. S. Kupper, "Antigen-independent processes in antigen-specific immunity.", J. Immunol., **150**, 1172 (1993) and P. L. Chisholm et al., "Monoclonal antibodies to the integrin  $\alpha_4$  subunit inhibit the murine contact hypersensitivity response." Eur. J. Immunol., **23**, 682 (1993)); viii) acute neurotoxic nephritis (see M. S. Mulligan et al., "Requirements for leukocyte adhesion molecules in nephrotoxic nephritis.", J. Clin. Invest., **91**, 577 (1993)); ix) tumor metastasis (for examples, see M. Edward, "Integrins and other adhesion molecules involved in melanocytic tumor progression.", Curr. Opin. Oncol., **7**, 185 (1995)); x) experimental autoimmune thyroiditis (see R. W. McMurray et al., "The role of  $\alpha_4$  integrin and intercellular adhesion molecule-1 (ICAM-1) in murine experimental autoimmune thyroiditis." Autoimmunity, **23**, 9 (1996); and xi) ischemic tissue damage following arterial occlusion in rats (see F. Squadrito et al., "Leukocyte integrin very late antigen-4/vascular cell adhesion molecule-1 adhesion pathway in splanchnic artery occlusion shock." Eur. J. Pharmacol., **318**, 153 (1996)). The primary mechanism of action of such antibodies appears to be the inhibition of lymphocyte and monocyte interactions with CAMs associated with components of the extracellular matrix, thereby limiting leukocyte migration to extravascular sites of injury or inflammation and/or limiting the priming and/or activation of leukocytes.

There is additional evidence supporting a possible role for VLA-4 interactions in other diseases, including rheumatoid arthritis; various melanomas, carcinomas, and sarcomas; inflammatory lung disorders; atherosclerotic plaque formation; restenosis; and circulatory



shock (for examples, see A. A. Postigo et al., "The  $\alpha_4\beta_1$ /VCAM-1 adhesion pathway in physiology and disease.", Res. Immunol., 144, 723 (1994) and J.-X. Gao and A. C. Issekutz, "Expression of VCAM-1 and VLA-4 dependent T-lymphocyte adhesion to dermal fibroblasts stimulated with proinflammatory cytokines." Immunol. 89, 375 (1996)).

At present, there is a humanized monoclonal antibody (Antegren® Athena Neurosciences/Elan ) against VLA-4 in clinical development for the treatment of "flares" associated with multiple sclerosis and a humanized monoclonal antibody (ACT-1® LeukoSite) against  $\alpha_4\beta_7$  in clinical development for the treatment of inflammatory bowel disease. Several peptidyl antagonists of VLA-4 have been described (D. Y. Jackson et al., "Potent  $\alpha_4\beta_1$  peptide antagonists as potential anti-inflammatory agents", J. Med. Chem., 40, 3359 (1997); H. N. Shroff et al., "Small peptide inhibitors of  $\alpha_4\beta_7$  mediated MadCAM-1 adhesion to lymphocytes", Bioorg. Med. Chem. Lett., 6, 2495 (1996); US 5,510,332, WO97/03094, WO97/02289, WO96/40781, WO96/22966, WO96/20216, WO96/01644, WO96/06108, WO95/15973). There is one report of nonpeptidyl inhibitors of the ligands for  $\alpha_4$ -integrins (WO96/31206). There still remains a need for low molecular weight, specific inhibitors of VLA-4- and  $\alpha_4\beta_7$ -dependent cell adhesion that have improved pharmacokinetic and pharmacodynamic properties such as oral bioavailability and significant duration of action. Such compounds would prove to be useful for the treatment, prevention or suppression of various pathologies mediated by VLA-4 and  $\alpha_4\beta_7$  binding and cell adhesion and activation.


#### SUMMARY OF THE INVENTION

The compounds of the present invention are antagonists of the VLA-4 integrin ("very late antigen-4"; CD49d/CD29; or  $\alpha_4\beta_1$ ) and/or the  $\alpha_4\beta_7$  integrin (LPAM-1 and  $\alpha_4\beta_7$ ), thereby blocking the binding of VLA-4 to its various ligands, such as VCAM-1 and regions of fibronectin and/or  $\alpha_4\beta_7$  to its various ligands, such as MadCAM-1, VCAM-1 and fibronectin. Thus, these antagonists are useful in inhibiting cell adhesion processes including cell activation, migration,

proliferation and differentiation. These antagonists are useful in the treatment, prevention and suppression of diseases mediated by VLA-4 and/or  $\alpha 4\beta 7$  binding and cell adhesion and activation, such as multiple sclerosis, asthma, allergic rhinitis, allergic conjunctivitis, inflammatory lung diseases, rheumatoid arthritis, septic arthritis, type I diabetes, organ transplantation, restenosis, autologous bone marrow transplantation, inflammatory sequelae of viral infections, myocarditis, inflammatory bowel disease including ulcerative colitis and Crohn's disease, certain types of toxic and immune-based nephritis, contact dermal hypersensitivity, psoriasis, tumor metastasis, and atherosclerosis.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention provides novel compounds of Formula I



I

or a pharmaceutically acceptable salt thereof wherein:

20

- R<sup>1</sup> is 1) C<sub>1</sub>-10alkyl,  
2) C<sub>2</sub>-10alkenyl,  
3) C<sub>2</sub>-10alkynyl,  
4) Cy,  
5) Cy-C<sub>1</sub>-10alkyl,  
6) Cy-C<sub>2</sub>-10alkenyl,  
7) Cy-C<sub>2</sub>-10alkynyl,

25

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from R<sup>a</sup>; and Cy is optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

30

R<sup>2</sup> and R<sup>3</sup> are independently

- 1) hydrogen, or
- 2) a group selected from R<sup>1</sup>; or

R<sup>2</sup> and R<sup>3</sup> together with the atoms to which they are attached form a ring of 4 to 7 members containing 0-2 additional heteroatoms

- 5 independently selected from oxygen, sulfur and nitrogen, wherein said ring may be isolated or benzo-fused, and optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

R<sup>4</sup> and R<sup>7</sup> are independently selected from the group consisting of

- 10
- 1) hydrogen,
  - 2) C<sub>1-10</sub>alkyl,
  - 3) C<sub>2-10</sub>alkenyl,
  - 4) C<sub>2-10</sub>alkynyl,
  - 5) aryl,
  - 15 6) aryl C<sub>1-10</sub>alkyl,
  - 7) heteroaryl, and
  - 8) heteroaryl C<sub>1-10</sub>alkyl,

wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents independently selected from R<sup>a</sup>, and aryl and

- 20 heteroaryl are optionally substituted with one to four substituents independently selected from R<sup>b</sup>; or

R<sup>3</sup>, R<sup>4</sup> and the carbon to which they are attached form a 3-7 membered ring optionally containing 0-2 heteroatoms selected from N, O and S;

25

- R<sup>5</sup> is
- 1) hydrogen,
  - 2) C<sub>1-10</sub>alkyl optionally substituted with one to four substituents independently selected from R<sup>a</sup>, or
  - 3) Cy optionally substituted with one to four substituents independently selected from R<sup>b</sup>,
- 30

- R<sup>6</sup> is
- 1) Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>1-10</sub>alkyl,
  - 2) Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>2-10</sub>alkenyl,
  - 3) Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>2-10</sub>alkynyl,

wherein Ar<sup>1</sup> and Ar<sup>2</sup> are independently aryl or heteroaryl each of which is optionally substituted with one to four substituents independently selected from R<sup>b</sup>; alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents independently  
 5 selected from R<sup>a</sup>;

- R<sup>a</sup> is
- 1) Cy
  - 2) -OR<sup>d</sup>,
  - 3) -NO<sub>2</sub>,
  - 10 4) halogen
  - 5) -S(O)<sub>m</sub>R<sup>d</sup>,
  - 6) -SR<sup>d</sup>,
  - 7) -S(O)<sub>2</sub>OR<sup>d</sup>,
  - 8) -S(O)<sub>m</sub>NR<sup>d</sup>Re,
  - 15 9) -NR<sup>d</sup>Re,
  - 10) -O(CR<sup>f</sup>R<sup>g</sup>)<sub>n</sub>NR<sup>d</sup>Re,
  - 11) -C(O)R<sup>d</sup>,
  - 12) -CO<sub>2</sub>R<sup>d</sup>,
  - 13) -CO<sub>2</sub>(CR<sup>f</sup>R<sup>g</sup>)<sub>n</sub>CONR<sup>d</sup>Re,
  - 20 14) -OC(O)R<sup>d</sup>,
  - 15) -CN,
  - 16) -C(O)NR<sup>d</sup>Re,
  - 17) -NR<sup>d</sup>C(O)R<sup>e</sup>,
  - 18) -OC(O)NR<sup>d</sup>Re,
  - 25 19) -NR<sup>d</sup>C(O)OR<sup>e</sup>,
  - 20) -NR<sup>d</sup>C(O)NR<sup>d</sup>Re,
  - 21) -CR<sup>d</sup>(N-OR<sup>e</sup>), or
  - 22) -CF<sub>3</sub>;

wherein Cy is optionally substituted with one to four substituents  
 30 independently selected from R<sup>c</sup>;

- R<sup>b</sup> is
- 1) a group selected from R<sup>a</sup>,
  - 2) C<sub>1</sub>-10 alkyl,

- 3) C<sub>2</sub>-10 alkenyl,
- 4) C<sub>2</sub>-10 alkynyl,
- 5) aryl C<sub>1</sub>-10alkyl,
- 6) heteroaryl C<sub>1</sub>-10 alkyl,

5 wherein alkyl, alkenyl, alkynyl, aryl, heteroaryl are optionally substituted with a group independently selected from R<sup>C</sup>;

- R<sup>C</sup> is
- 1) halogen,
  - 2) amino,
  - 10 3) carboxy,
  - 4) C<sub>1</sub>-4alkyl,
  - 5) C<sub>1</sub>-4alkoxy,
  - 6) aryl,
  - 7) aryl C<sub>1</sub>-4alkyl, or
  - 15 8) aryloxy.

R<sup>d</sup> and R<sup>e</sup> are independently selected from hydrogen, C<sub>1</sub>-10alkyl, C<sub>2</sub>-10alkenyl, C<sub>2</sub>-10alkynyl, Cy and Cy C<sub>1</sub>-10alkyl, wherein alkyl, alkenyl, alkynyl and Cy is optionally substituted with one to four  
20 substituents independently selected from R<sup>C</sup>; or  
R<sup>d</sup> and R<sup>e</sup> together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

25 R<sup>f</sup> and R<sup>g</sup> are independently selected from hydrogen, C<sub>1</sub>-10alkyl, Cy and Cy C<sub>1</sub>-10alkyl; or  
R<sup>f</sup> and R<sup>g</sup> together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

30

- R<sup>h</sup> is
- 1) hydrogen,
  - 2) C<sub>1</sub>-10alkyl,
  - 3) C<sub>2</sub>-10alkenyl,
  - 4) C<sub>2</sub>-10alkynyl,

- 5) cyano,
- 6) aryl,
- 7) aryl C<sub>1-10</sub>alkyl,
- 8) heteroaryl,
- 5 9) heteroaryl C<sub>1-10</sub>alkyl, or
- 10) -SO<sub>2</sub>R<sup>i</sup>;

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from R<sup>a</sup>; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

- R<sup>i</sup>
- 1) C<sub>1-10</sub>alkyl,
  - 2) C<sub>2-10</sub>alkenyl,
  - 3) C<sub>2-10</sub>alkynyl, or
  - 15 4) aryl;

wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from R<sup>c</sup>;

Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

m is an integer from 1 to 2;

n is an integer from 1 to 10;

- 25 X is 1) -C(O)OR<sup>d</sup>,
- 2) -P(O)(OR<sup>d</sup>)(OR<sup>e</sup>)
- 3) -P(O)(R<sup>d</sup>)(OR<sup>e</sup>)
- 4) -S(O)<sub>m</sub>OR<sup>d</sup>,
- 5) -C(O)NR<sup>d</sup>R<sup>h</sup>, or
- 30 6) -5-tetrazolyl;

- Y is
- 1) -C(O)-,
  - 2) -O-C(O)-,
  - 3) -NR<sup>e</sup>-C(O)-,

- 4) -S(O)<sub>2</sub>-,
- 5) -P(O)(OR<sup>i</sup>)
- 6) C(O)C(O).

5                    In one embodiment, R<sup>1</sup> is C<sub>1</sub>-10alkyl, aryl, aryl-C<sub>1</sub>-10alkyl, heteroaryl or heteroaryl-C<sub>1</sub>-10alkyl, wherein alkyl, aryl and heteroaryl are optionally substituted as provided for under formula I. In a preferred embodiment R<sup>1</sup> is phenyl optionally substituted with one to three groups selected from R<sup>b</sup>.


10                   In another embodiment, R<sup>2</sup> is hydrogen, C<sub>1</sub>-10 alkyl, Cy or Cy-C<sub>1</sub>-10 alkyl; or R<sup>2</sup>, R<sup>3</sup> together with the atoms to which they are attached form a ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen, wherein said ring may be isolated or benzo-fused, and optionally  
15 substituted with one to four substituents independently selected from R<sup>b</sup>. Preferably R<sup>2</sup>, R<sup>3</sup> together with the atoms to which they are attached for a ring of 5 to 6 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen, wherein said ring may be isolated or benzo-fused, and optionally substituted with one  
20 to four substituents independently selected from R<sup>b</sup>.

                    In another embodiment R<sup>4</sup> is H, C<sub>1</sub>-10alkyl, aryl, heteroaryl, aryl-C<sub>1</sub>-10alkyl or heteroaryl-C<sub>1</sub>-10alkyl. Preferably, R<sup>4</sup> is H or C<sub>1</sub>-10 alkyl.

                    In another embodiment R<sup>6</sup> is Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>1</sub>-10alkyl wherein  
25 Ar<sup>1</sup> and Ar<sup>2</sup> are optionally substituted with from 1 to 4 groups independently selected from R<sup>b</sup>. Preferably R<sup>6</sup> is Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>1</sub>-3alkyl wherein Ar<sup>1</sup> and Ar<sup>2</sup> are optionally substituted with from 1 to 4 groups independently selected from R<sup>b</sup>.

                    In another embodiment X is C(O)OR<sup>d</sup>.  
30                   In yet another embodiment Y is C(O) or S(O)<sub>2</sub>.

                    In a preferred embodiment of compounds of Formula I are compounds of formula Ia:



wherein

R<sup>1</sup> is 1) C<sub>1-10</sub>alkyl,

5           2) Cy, or

          3) Cy-C<sub>1-10</sub>alkyl,

wherein alkyl is optionally substituted with one to four substituents independently selected from R<sup>a</sup>; and Cy is optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

10   R<sup>4</sup> is 1) hydrogen, or

          2) C<sub>1-3</sub> alkyl;

Ar<sup>1</sup> and Ar<sup>2</sup> or independently aryl or heteroaryl each of which is optionally substituted with one to four substituents independently selected from R<sup>b</sup>; and

15   R<sup>b</sup> is as defined under formula I.

Representative compounds of formula I are as follows:

N-(3,4-dimethoxybenzenesulfonyl)-1,2,3,4-tetrahydro-isoquinoline-3(S)-carbonyl-(L)-biphenylalanine;

N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-biphenylalanine;

20   N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4-fluorophenyl)phenylalanine;

N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-thienyl)-phenylalanine;

25   N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-thienyl)-phenylalanine;

N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4'-trifluoromethyl-phenyl)-phenylalanine;

N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methoxy-phenyl)-phenylalanine;

30   N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-formyl-phenyl)-phenylalanine;



- N-(3-fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-thienyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2',6'-difluorophenyl)phenylalanine;  
5 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-hydroxymethylphenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4'-methylphenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-carboxyphenyl)phenylalanine;  
10 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methoxycarbonylphenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-formylphenyl)phenylalanine;  
15 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-aminophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylphenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-acetamidophenyl)phenylalanine;  
20 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-fluorophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-carboxyphenyl)phenylalanine;  
25 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-methoxycarbonylphenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2',4'-dichlorophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-formyl-3-thienyl)phenylalanine;  
30 N-(3,5-dichlorobenzenesulfonyl)-2(S)-methylprolyl-(L)-4-(4'-fluorophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-formylphenyl)phenylalanine;

- N-(3,5-dichlorobenzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-(hydroxymethyl)phenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-cyanophenyl)phenylalanine;
- 5 N-(benzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-formylphenyl)phenylalanine;  
N-(benzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-methoxyphenyl)phenylalanine;
- 10 N-(3-Fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4'-fluoro-2'-methoxyphenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;
- 15 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylthiophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(5-methyl-1,3,4-oxadiazol-2-yl-phenyl)phenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-methyl-5-
- 20 trifluoromethyl-benzoxazol-7-yl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-methyl-6-(5-trifluoromethyl-tetrazol-1-yl)-benzoxazol-4-yl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-methyl-5-(5-trifluoromethyl-tetrazol-1-yl)-benzoxazol-7-yl)phenylalanine;
- 25 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3-pyridyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-pyridyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(5-
- 30 pyrimidinyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-cyanophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-methyl-benzoxazol-4-yl)phenylalanine;

- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(6-acetamido-2-methyl-benzoxazol-4-yl)phenylalanine;  
N-(benzenesulfonyl)-(L)-prolyl-(L)-4-(2-pyridyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-3(S)-methylprolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
5 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(tetrazol-5-yl)phenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(2-methyl-tetrazol-5-yl)phenyl)phenylalanine;  
10 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(3-methyl-tetrazol-5-yl)phenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-aminocarbonylphenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylaminocarbonylphenyl)phenylalanine;  
15 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(benzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-carboxyphenyl)phenylalanine;  
20 N-(3-bromobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(benzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-( $\alpha$ -toluenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(phenylacetyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;;  
25 N-(3-pyridinesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(2-thienylsulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(benzylaminocarbonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(3-phenylpropionyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
30 N-((5-methyl-3,4-thiadiazol-2-yl)sulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-((benzthiazol-2-yl)sulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;

- N-((1-methyl-imidazol-4-yl)sulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylsulfinylphenyl)phenylalanine; and  
5 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylsulfonylphenyl)phenylalanine.

"Alkyl", as well as other groups having the prefix "alk", such as alkoxy, alkanoyl, means carbon chains which may be linear or  
10 branched or combinations thereof. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, sec- and tert-butyl, pentyl, hexyl, heptyl, octyl, nonyl, and the like.

"Alkenyl" means carbon chains which contain at least one carbon-carbon double bond, and which may be linear or branched or  
15 combinations thereof. Examples of alkenyl include vinyl, allyl, isopropenyl, pentenyl, hexenyl, heptenyl, 1-propenyl, 2-butenyl, 2-methyl-2-butenyl, and the like.

"Alkynyl" means carbon chains which contain at least one carbon-carbon triple bond, and which may be linear or branched or  
20 combinations thereof. Examples of alkynyl include ethynyl, propargyl, 3-methyl-1-pentynyl, 2-heptynyl and the like.

"Cycloalkyl" means mono- or bicyclic saturated carbocyclic rings, each of which having from 3 to 10 carbon atoms. The term also includes monocyclic ring fused to an aryl group in which the point of  
25 attachment is on the non-aromatic portion. Examples of cycloalkyl include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, tetrahydronaphthyl, decahydronaphthyl, indanyl, and the like.

"Aryl" means mono- or bicyclic aromatic rings containing only carbon atoms. The term also includes aryl group fused to a  
30 monocyclic cycloalkyl or monocyclic heterocyclyl group in which the point of attachment is on the aromatic portion. Examples of aryl include phenyl, naphthyl, indanyl, indenyl, tetrahydronaphthyl, 2,3-dihydrobenzofuranyl, benzopyranyl, 1,4-benzodioxanyl, and the like.

"Heteroaryl" means a mono- or bicyclic aromatic ring containing at least one heteroatom selected from N, O and S, with each ring containing 5 to 6 atoms. Examples of heteroaryl include pyrrolyl, isoxazolyl, isothiazolyl, pyrazolyl, pyridyl, oxazolyl, oxadiazolyl, thiadiazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, furanyl, triazinyl, thienyl, pyrimidyl, pyridazinyl, pyrazinyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, benzothiophenyl, furo(2,3-b)pyridyl, quinolyl, indolyl, isoquinolyl, and the like.

"Heterocyclyl" means mono- or bicyclic saturated rings containing at least one heteroatom selected from N, S and O, each of said ring having from 3 to 10 atoms. The term also includes monocyclic heterocycle fused to an aryl or heteroaryl group in which the point of attachment is on the non-aromatic portion. Examples of "heterocyclyl" include pyrrolidinyl, piperidinyl, piperazinyl, imidazolidinyl, 2,3-dihydrofuro(2,3-b)pyridyl, benzoxazinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, dihydroindolyl, and the like.

"Halogen" includes fluorine, chlorine, bromine and iodine.

#### 20 Optical Isomers - Diastereomers - Geometric Isomers

Compounds of Formula I contain one or more asymmetric centers and can thus occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. The present invention is meant to comprehend all such isomeric forms of the compounds of Formula I.

Some of the compounds described herein contain olefinic double bonds, and unless specified otherwise, are meant to include both E and Z geometric isomers.

#### 30 Salts

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Salts derived from inorganic bases include aluminum, ammonium,

calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc, and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium, and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like.

When the compound of the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pantoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid, and the like. Particularly preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

It will be understood that, as used herein, references to the compounds of Formula I are meant to also include the pharmaceutically acceptable salts.

### Utilities

The ability of the compounds of Formula I to antagonize the actions of VLA-4 and/or  $\alpha 4\beta 7$  integrin makes them useful for preventing or reversing the symptoms, disorders or diseases induced by the binding of VLA-4 and/or  $\alpha 4\beta 7$  to their various respective ligands. Thus, these antagonists will inhibit cell adhesion processes including cell activation, migration, proliferation and differentiation. Accordingly,

another aspect of the present invention provides a method for the treatment (including prevention, alleviation, amelioration or suppression) of diseases or disorders or symptoms mediated by VLA-4 and/or  $\alpha 4\beta 7$  binding and cell adhesion and activation, which comprises  
5 administering to a mammal an effective amount of a compound of Formula I. Such diseases, disorders, conditions or symptoms are for example (1) multiple sclerosis, (2) asthma, (3) allergic rhinitis, (4) allergic conjunctivitis, (5) inflammatory lung diseases, (6) rheumatoid arthritis, (7) septic arthritis, (8) type I diabetes, (9) organ  
10 transplantation rejection, (10) restenosis, (11) autologous bone marrow transplantation, (12) inflammatory sequelae of viral infections, (13) myocarditis, (14) inflammatory bowel disease including ulcerative colitis and Crohn's disease, (15) certain types of toxic and immune-based nephritis, (16) contact dermal hypersensitivity, (17) psoriasis,  
15 (18) tumor metastasis, and (19) atherosclerosis.

#### Dose Ranges

The magnitude of prophylactic or therapeutic dose of a compound of Formula I will, of course, vary with the nature of the  
20 severity of the condition to be treated and with the particular compound of Formula I and its route of administration. It will also vary according to the age, weight and response of the individual patient. In general, the daily dose range lie within the range of from about 0.001 mg to about 100 mg per kg body weight of a mammal, preferably 0.01 mg to about  
25 50 mg per kg, and most preferably 0.1 to 10 mg per kg, in single or divided doses. On the other hand, it may be necessary to use dosages outside these limits in some cases.

For use where a composition for intravenous administration is employed, a suitable dosage range is from about 0.001 mg to about 25  
30 mg (preferably from 0.01 mg to about 1 mg) of a compound of Formula I per kg of body weight per day and for cytoprotective use from about 0.1 mg to about 100 mg (preferably from about 1 mg to about 100 mg and more preferably from about 1 mg to about 10 mg) of a compound of Formula I per kg of body weight per day.

In the case where an oral composition is employed, a suitable dosage range is, e.g. from about 0.01 mg to about 100 mg of a compound of Formula I per kg of body weight per day, preferably from about 0.1 mg to about 10 mg per kg and for cytoprotective use  
5 from 0.1 mg to about 100 mg (preferably from about 1 mg to about 100 mg and more preferably from about 10 mg to about 100 mg) of a compound of Formula I per kg of body weight per day.

For the treatment of diseases of the eye, ophthalmic preparations for ocular administration comprising 0.001-1% by weight  
10 solutions or suspensions of the compounds of Formula I in an acceptable ophthalmic formulation may be used.

#### Pharmaceutical Compositions

Another aspect of the present invention provides  
15 pharmaceutical compositions which comprises a compound of Formula I and a pharmaceutically acceptable carrier. The term "composition", as in pharmaceutical composition, is intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) (pharmaceutically acceptable excipients) that make up the carrier, as  
20 well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the  
25 present invention encompass any composition made by admixing a compound of Formula I, additional active ingredient(s), and pharmaceutically acceptable excipients.

Any suitable route of administration may be employed for providing a mammal, especially a human with an effective dosage of a  
30 compound of the present invention. For example, oral, rectal, topical, parenteral, ocular, pulmonary, nasal, and the like may be employed. Dosage forms include tablets, troches, dispersions, suspensions, solutions, capsules, creams, ointments, aerosols, and the like.



The pharmaceutical compositions of the present invention comprise a compound of Formula I as an active ingredient or a pharmaceutically acceptable salt thereof, and may also contain a pharmaceutically acceptable carrier and optionally other therapeutic ingredients. The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic bases or acids and organic bases or acids.

The compositions include compositions suitable for oral, rectal, topical, parenteral (including subcutaneous, intramuscular, and intravenous), ocular (ophthalmic), pulmonary (nasal or buccal inhalation), or nasal administration, although the most suitable route in any given case will depend on the nature and severity of the conditions being treated and on the nature of the active ingredient. They may be conveniently presented in unit dosage form and prepared by any of the methods well-known in the art of pharmacy.

For administration by inhalation, the compounds of the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or nebulisers. The compounds may also be delivered as powders which may be formulated and the powder composition may be inhaled with the aid of an insufflation powder inhaler device. The preferred delivery system for inhalation is a metered dose inhalation (MDI) aerosol, which may be formulated as a suspension or solution of a compound of Formula I in suitable propellants, such as fluorocarbons or hydrocarbons.

Suitable topical formulations of a compound of formula I include transdermal devices, aerosols, creams, ointments, lotions, dusting powders, and the like.

In practical use, the compounds of Formula I can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., oral or parenteral (including intravenous). In preparing the compositions for oral dosage form, any of the usual pharmaceutical

media may be employed, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like in the case of oral liquid preparations, such as, for example, suspensions, elixirs and solutions; or carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents and the like in the case of oral solid preparations such as, for example, powders, capsules and tablets, with the solid oral preparations being preferred over the liquid preparations. Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit form in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be coated by standard aqueous or nonaqueous techniques.

In addition to the common dosage forms set out above, the compounds of Formula I may also be administered by controlled release means and/or delivery devices such as those described in U.S. Patent Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; 3,630,200 and 4,008,719.

Pharmaceutical compositions of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient, as a powder or granules or as a solution or a suspension in an aqueous liquid, a non-aqueous liquid, an oil-in-water emulsion or a water-in-oil liquid emulsion. Such compositions may be prepared by any of the methods of pharmacy but all methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product into the desired presentation. For example, a tablet may be prepared by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine, the active ingredient in a free-flowing form such as powder or granules, optionally mixed with a binder, lubricant, inert diluent,

surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. Desirably, each tablet contains from about 1 mg to about 500 mg of the active ingredient and each  
 5 cachet or capsule contains from about 1 to about 500 mg of the active ingredient.

The following are examples of representative pharmaceutical dosage forms for the compounds of Formula I:

10	<u>Injectable Suspension (I.M.)</u>	<u>mg/mL</u>
	Compound of Formula I 10	
	Methylcellulose	5.0
	Tween 80	0.5
	Benzyl alcohol	9.0
15	Benzalkonium chloride	1.0
	Water for injection to a total volume of 1 mL	

	<u>Tablet</u>	<u>mg/tablet</u>
	Compound of Formula I 25	
20	Microcrystalline Cellulose	415
	Povidone	14.0
	Pregelatinized Starch	3.5
	Magnesium Stearate	<u>2.5</u>
		500

25	<u>Capsule</u>	<u>mg/capsule</u>
	Compound of Formula I 25	
	Lactose Powder	573.5
	Magnesium Stearate	<u>1.5</u>
30		600

	<u>Aerosol</u>	<u>Per canister</u>
	Compound of Formula I 24 mg	
	Lecithin, NF Liquid Conc.	1.2 mg

Trichlorofluoromethane, NF 4.025 g

Dichlorodifluoromethane, NF 12.15 g

### Combination Therapy

- 5                   Compounds of Formula I may be used in combination with other drugs that are used in the treatment/prevention/suppression or amelioration of the diseases or conditions for which compounds of Formula I are useful. Such other drugs may be administered, by a route and in an amount commonly used therefor, contemporaneously or
- 10 sequentially with a compound of Formula I. When a compound of Formula I is used contemporaneously with one or more other drugs, a pharmaceutical composition containing such other drugs in addition to the compound of Formula I is preferred. Accordingly, the pharmaceutical compositions of the present invention include those that
- 15 also contain one or more other active ingredients, in addition to a compound of Formula I. Examples of other active ingredients that may be combined with a compound of Formula I, either administered separately or in the same pharmaceutical compositions, include, but are not limited to:
- 20 (a) other VLA-4 antagonists such as those described in US 5,510,332, WO97/03094, WO97/02289, WO96/40781, WO96/22966, WO96/20216, WO96/01644, WO96/06108, WO95/15973 and WO96/31206; (b) steroids such as beclomethasone, methylprednisolone, betamethasone, prednisone, dexamethasone, and hydrocortisone; (c)
- 25 immunosuppressants such as cyclosporin, tacrolimus, rapamycin and other FK-506 type immunosuppressants; (d) antihistamines (H1-histamine antagonists) such as bromopheniramine, chlorpheniramine, dexchlorpheniramine, triprolidine, clemastine, diphenhydramine, diphenylpyraline, tripelemamine, hydroxyzine, methdilazine,
- 30 promethazine, trimeprazine, azatadine, cyproheptadine, antazoline, pheniramine pyrilamine, astemizole, terfenadine, loratadine, cetirizine, fexofenadine, descarboethoxyloratadine, and the like; (e) non-steroidal anti-asthmatics such as  $\beta$ 2-agonists (terbutaline, metaproterenol, fenoterol, isoetharine, albuterol, bitolterol, and pirbuterol),


theophylline, cromolyn sodium, atropine, ipratropium bromide, leukotriene antagonists (zafirlukast, montelukast, pranlukast, iralukast, pobilukast, SKB-106,203), leukotriene biosynthesis inhibitors (zileuton, BAY-1005); (f) non-steroidal antiinflammatory agents (NSAIDs) such as propionic acid derivatives (alminoprofen, benoxaprofen, bucloxic acid, carprofen, fenbufen, fenoprofen, fluprofen, flurbiprofen, ibuprofen, indoprofen, ketoprofen, miroprofen, naproxen, oxaprozin, pirprofen, pranoprofen, suprofen, tiaprofenic acid, and tioxaprofen), acetic acid derivatives (indomethacin, acemetacin, alclofenac, clidanac, diclofenac, fenclofenac, fenclozic acid, fentiazac, furofenac, ibufenac, isoxepac, oxpinac, sulindac, tiopinac, tolmetin, zidometacin, and zomepirac), fenamic acid derivatives (flufenamic acid, meclofenamic acid, mefenamic acid, niflumic acid and tolfenamic acid), biphenylcarboxylic acid derivatives (diflunisal and flufenisal), oxicams (isoxicam, piroxicam, sudoxicam and tenoxicam), salicylates (acetyl salicylic acid, sulfasalazine) and the pyrazolones (apazone, bezpiperylon, feprazone, mofebutazone, oxyphenbutazone, phenylbutazone); (g) cyclooxygenase-2 (COX-2) inhibitors; (h) inhibitors of phosphodiesterase type IV (PDE-IV); (i) antagonists of the chemokine receptors, especially CCR-1, CCR-2, and CCR-3; (j) cholesterol lowering agents such as HMG-CoA reductase inhibitors (lovastatin, simvastatin and pravastatin, fluvastatin, atorvastatin, and other statins), sequestrants (cholestyramine and colestipol), nicotinic acid, fenofibric acid derivatives (gemfibrozil, clofibrat, fenofibrate and benzaifibrate), and probucol; (k) anti-diabetic agents such as insulin, sulfonylureas, biguanides (metformin),  $\alpha$ -glucosidase inhibitors (acarbose) and glitazones (troglitazone and pioglitazone); (l) preparations of interferon beta (interferon beta-1a, interferon beta-1b); (m) anticholinergic agents such as muscarinic antagonists (ipratropium bromide); (n) other compounds such as 5-aminosalicylic acid and prodrugs thereof, antimetabolites such as azathioprine and 6-mercaptopurine, and cytotoxic cancer chemotherapeutic agents.

The weight ratio of the compound of the Formula I to the second active ingredient may be varied and will depend upon the

effective dose of each ingredient. Generally, an effective dose of each will be used. Thus, for example, when a compound of the Formula I is combined with an NSAID the weight ratio of the compound of the Formula I to the NSAID will generally range from about 1000:1 to about 1:1000, preferably about 200:1 to about 1:200. Combinations of a compound of the Formula I and other active ingredients will generally also be within the aforementioned range, but in each case, an effective dose of each active ingredient should be used.

Compounds of the present invention may be prepared by procedures illustrated in the accompanying schemes. In the first method (Scheme 1), a resin-based synthetic strategy is outlined where the resin employed is represented by the ball (●). An N-Fmoc-protected amino acid derivative A (Fmoc = fluorenylmethoxycarbonyl) is loaded on to the appropriate hydroxyl-containing resin using dicyclohexylcarbodiimide (DCC) and 1-hydroxybenzotriazole (HOBt) in dimethylformamide (DMF) to give B. The Fmoc protecting group is removed with piperidine in DMF to yield free amine C. The next Fmoc-protected amino acid derivative D is coupled to C employing standard peptide (in this instance, 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU), HOBt, and N,N-diisopropylethylamine (DIEA) in DMF) to yield dipeptide E. The Fmoc group is removed with piperidine in DMF to yield the free amine F. An acid chloride or isocyanate derivative is reacted with F in the presence of DIEA to yield G. The final product is removed from the resin with strong acid (in this instance, trifluoroacetic acid (TFA) in the presence of thioanisole and ethanedithiol) to yield compounds of the present invention H.


Scheme 1.



- In the second method (Scheme 2), standard solution phase synthetic methodology is outlined. An N-Boc-protected amino acid derivative **A** (Boc = tert-butyloxycarbonyl) is treated with tert-butyl 2,2,2-trichloroacetimidate in the presence of boron trifluoride etherate to yield tert-butyl ester **B** which is subsequently coupled to Cbz-protected amino acid derivative **C** (Cbz = carbobenzyloxy) in the presence of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC), HOBT, and N-methylmorpholine (NMM) in methylene chloride ( $\text{CH}_2\text{Cl}_2$ ) to yield dipeptide **D**. Catalytic hydrogenation of **D** in the presence of a palladium-on-carbon (Pd/C) catalyst yields **E**. Reaction of **E** with an acid chloride or isocyanate in the presence of DIEA and 4-dimethylaminopyridine (DMAP) yields **F**.

which is subsequently reacted with strong acid (TFA) to yield the desired product G.

Scheme 2.



5


In the third method (Scheme 3), a late stage intermediate aryl bromide or iodide is coupled to an appropriately substituted aryl or heteroaryl boronic acid to give a subset of compounds of the present invention ( $\text{R}^6$  = biaryl-substituted alkyl,  $\text{R}^7$  = hydrogen). For example, amino acid methyl ester A is reacted with an acid chloride or isocyanate in the presence of DIEA to yield B. Basic hydrolysis of the methyl ester yields amino acid derivative C. N-Boc-4-iodo- or 4-bromophenylalanine D is reacted with *tert*-butyl 2,2,2-trichloroacetimidate in the presence of boron trifluoride etherate in methylene chloride-cyclohexane to yield *tert*-butyl ester E which is subsequently coupled with C in the presence of EDC, HOBT and NMM to yield 4-iodo- or 4-bromo-phenylalanine dipeptide F. Substituted aryl or heteroaryl boronic acids are coupled to F in the presence of a palladium(0) reagent, such as tetrakis(triphenylphosphine)palladium under Suzuki conditions (N. Miyaura *et al.*, *Synth. Commun.*, **1981**, *11*, 513-519) to yield G. The *tert*-butyl ester is then removed by treatment with strong

20



- acid (TFA) to yield the desired product **H**. If the aryl or heteroaryl boronic acid is not commercially available, but the corresponding bromide or iodide is, then the bromide or iodide can be converted into the desired boronic acid by treatment with an alkyllithium reagent in tetrahydrofuran at low temperature followed by addition of trimethyl or triisopropyl borate. Hydrolysis to the boronic acid can be effected by treatment of the intermediate with aqueous base and then acid.

Scheme 3.




10

- Alternatively, the aryl coupling reaction may be performed by application of Stille-type carbon-carbon bond forming conditions (Scheme 4). (A.M. Echavarren and J.K. Stille, *J. Am. Chem. Soc.* **1987**, *109*, 5478-5486). The aryl bromide or iodide intermediate **A** is converted into its trimethyltin derivative **B** using hexamethylditin in the

15

- presence of palladium(0) and lithium chloride and then reacted with an appropriately substituted aryl or heteroaryl bromide, iodide, or triflate in the presence of a palladium reagent, such as tetrakis(triphenylphosphine)palladium(0) or
- 5 tris(dibenzylideneacetone)dipalladium(0), in a suitable solvent, such as toluene, dioxane, DMF, or 1-methyl-2-pyrrolidinone, to give intermediate **C**. The tert-butyl ester is then removed by treatment with strong acid (TFA) to yield the desired product **D**.

Scheme 4.




10

- Compounds wherein the middle ring is heteroaryl (**E**) may be prepared (Scheme 5) in a similar fashion starting from the appropriate heteroaryl bromide or iodide **C** using Suzuki-type
- 15 conditions as depicted in Scheme 3 or from the corresponding heteroaryl trimethyltin intermediate **D** using Stille-type conditions as depicted in Scheme 4. The requisite heteroaryl halides **C** may be prepared via conventional electrophilic halogenation of the N-Boc-heteroaryl-alanine tert-butyl ester intermediate **B**. **B** may be prepared

from the known aliphatic iodo intermediate **A** in carbon-carbon bond formation using zinc/copper couple and palladium(II) (M.J. Dunn *et al.*, *SYNLETT* 1993, 499-500).

Scheme 5.



5

### General procedure for the solid-phase synthesis of compounds of Formula 1.

#### 10 Step A. Loading of N-Fmoc-amino acid derivatives onto resins.

N-Fmoc-amino acids were loaded on either Wang® (Calbiochem-Novabiochem Corp.) or Chloro (2-chlorotrityl) resin.

Wang® resin, typically 0.3 mmol, was washed with dimethylformamide three times. A solution of N-Fmoc-amino acid (0.3 mmol) in dimethylformamide (3 mL) was transferred to the pre-swollen Wang® resin. Dicyclohexylcarbodiimide (0.3 mmol) and 1-N-

5 hydroxybenzotriazole (0.3 mmol) was added and the mixture gently swirled for 2 hours. Following filtration, the resin was sequentially washed with dimethylformamide (3 times) and dichloromethane (3 times). The amino acid substitution value obtained after vacuum drying typically ranged between 0.07 to 0.1 mmol.

10 Alternatively, Chloro (2-chlorotriyl) resin, typically 0.2 mmol, was pre-swollen in dimethylformamide. A solution of N-Fmoc-amino acid (0.2 mmol) in dimethylformamide (3 ml) was added to the resin, followed by the addition of N,N-diisopropylethylamine (0.4 mmol). The resin was gently stirred for 2 hours, filtered and washed  
15 sequentially with dimethylformamide (3 times) and dichloromethane (3 times). The resin was finally washed with 10% methanol in dichloromethane and vacuum dried. The amino acid substitution value obtained after vacuum drying typically ranged between 0.05 to 0.1 mmol.

20

Step B. Deprotection of the N-Fmoc group.

The N-Fmoc protecting group was removed from the resin from Step A by treatment with 20% piperidine in dimethylformamide for 30 minutes. Following filtration, the resin was washed sequentially  
25 with dimethylformamide (3 times), dichloromethane (1 time) and dimethylformamide (2 times) and used in the subsequent reaction.

Step C. Coupling of the next N-Fmoc-amino acid derivative

A solution of the next desired N-Fmoc-amino acid  
30 derivative (0.4 mmol) in dimethylformamide (2 mL) was mixed with 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (0.4 mmol), 1-hydroxybenzotriazole (0.4 mmol) and diisopropylethylamine (0.6 mmol). This solution was transferred to resin from Step B and typically allowed to react for 2 hours. Couplings

were monitored by ninhydrin reaction. The coupling mixture was filtered and the resin washed with dimethylformamide (3 times) and used in the subsequent reaction.

5 Step D. Deprotection of the N-Fmoc group.

The N-Fmoc protecting group was removed from the resin from Step C by the procedure described in Step B and used in the subsequent reaction.

10 Step E. Acylation (or sulfonylation) of the terminal amino group.

The desired N-terminal capping reagent (sulfonyl chloride or other acyl chloride) (0.4 mol) was dissolved in dimethylformamide (2 ml), mixed with N,N-diisopropylethylamine (0.8 mmol) and added to the resin from Step D. After approximately two hours, the resin was  
15 sequentially washed with dimethylformamide (3 times) and dichloromethane (3 times).

Step F. Cleavage of the desired products from the resins.

The final desired products were cleaved from the resins  
20 from Step E by gently stirring with a solution of trifluoroacetic acid:thioanisole:ethanedithiol (95:2.5:2.5); 3 hours for Wang® resin and 30 minutes for the Chloro (2-chlorotriyl) resin. Following filtration, the solvents were removed by evaporation and the residue dissolved in acetonitrile (3 mL). Insoluble material was removed by filtration. The  
25 final products were purified by reverse phase chromatography with a linear gradient of buffer A (0.1% trifluoroacetic acid in water) and buffer B (0.1% trifluoroacetic acid in acetonitrile) and isolated by lyophilization. Molecular ions were obtained by electrospray ionization mass spectrometry or matrix-assisted laser desorption ionization time-  
30 of-flight mass spectrometry to confirm the structure of each peptide.

The following compounds were prepared by the above general procedures using the appropriate amino acid derivatives and acylor sulfonyl chloride:



Example	Compound Name	MS *
(1)	N-(3,4-dimethoxybenzenesulfonyl)-1,2,3,4-tetrahydro-isoquinoline-3(S)-carbonyl-(L)-biphenylalanine	601
(2)	N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-biphenylalanine	548

\* m/e, M+1

### EXAMPLE 3

#### 5 N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4-fluorophenyl)phenylalanine

Step A: 4-Iodo-(L)-phenylalanine, tert-butyl ester hydrochloride.

To a suspension of N-Boc-4-iodo-(L)-phenylalanine (1.0 g, 2.56 mmol) in methylene chloride (7 mL) and cyclohexane (14 mL)  
 10 were added t-butyl trichloroacetimidate (0.48 mL, 2.68 mmol) and boron trifluoride-etherate (48  $\mu$ L). The reaction mixture was stirred for 5 hours at room temperature under a nitrogen atmosphere and then treated a second time with the same amounts of t-butyl trichloroacetimidate and boron trifluoride-etherate as above. After  
 15 stirring overnight, a third addition was made, and the mixture was stirred a further 3 hours. The mixture was then filtered and the filtrate evaporated. The product was obtained pure by silica gel chromatography eluting with 10% diethyl ether in hexane; yield 650 mg. The product was treated with 1M HCl in ethyl acetate (7.3 mL) for  
 20 18 hours at room temperature. The mixture was evaporated and coevaporated several times with diethyl ether to afford the title compound; yield 522 mg.  
 400 MHz  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.42 (s, 9H); 3.13 (d, 2H); 4.18 (t, 1H); 7.09 (d, 2H); 7.75 (d, 2H).

25

Step B: N-(3,5-Dichlorobenzenesulfonyl)-(L)-proline

To a mixture of (L)-proline methyl ester hydrochloride (838 mg, 5.06 mmol) in methylene chloride (25 mL) at 0°C were added N,N-

diisopropylethylamine (2.64 mL, 15.2 mmol) and a solution of 3,5-dichlorobenzenesulfonyl chloride (1.49 g, 6.07 mmol) in methylene chloride (5 mL). The cooling bath was removed, and the mixture was stirred overnight at room temperature. It was then diluted with

5 methylene chloride, washed with 1N hydrochloric acid, saturated NaHCO<sub>3</sub>, saturated brine solution, dried (Na<sub>2</sub>SO<sub>4</sub>), and evaporated.

The methyl ester was obtained pure by silica gel chromatography eluting with 10% acetone in hexane; yield 1.49 g. It was then taken up in ethanol (50 mL) and treated with 0.2 N sodium hydroxide (26.6 mL) for 1.5 hours at room temperature. The mixture was acidified with glacial acetic acid, concentrated, the residue taken up in methylene chloride, washed with water, saturated brine solution, dried (Na<sub>2</sub>SO<sub>4</sub>), and evaporated to give the title compound; yield 1.4 g.

10 400 MHz <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 1.80-2.15 (m, 4H); 3.35-4.45 (m, 2H); 4.30 (dd, 1H); 7.76 (m, 1H); 7.83 (m, 2H).

Step C: N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-iodophenylalanine, tert-butyl ester.

To a solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-proline (386 mg, 1.19 mmol) in methylene chloride (23 mL) were added 1-hydroxybenzotriazole (241 mg, 1.79 mmol), N-methylmorpholine (0.33 mL, 2.98 mmol), and 4-iodo-(L)-phenylalanine tert-butyl ester hydrochloride (458 mg, 1.19 mmol). After cooling in an ice-bath for 5 minutes, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (274 mg, 1.43 mmol) was added. After 15 minutes, the cooling bath was removed, and the mixture was stirred overnight under a nitrogen atmosphere. The mixture was diluted with methylene chloride, washed with water, 1N HCl, saturated NaHCO<sub>3</sub> solution, saturated brine solution, dried (MgSO<sub>4</sub>), and evaporated. Silica gel chromatography eluting with 20% ethyl acetate in hexane afforded pure title compound; yield 651 mg (84%).

MS: m/e 653 (M + 1)



400 MHz  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.45 (s, 9H); 1.65-1.85 (m, 4H); 3.0 (dd, 1H); 3.13 (dd, 1H); 3.45 (m, 1H); 4.20 (m, 1H); 4.55 (dd, 1H); 7.05 (d, 2H); 7.64 (d, 2H); 7.80 (s, 3H).

5 Step D: N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4-fluorophenyl)phenylalanine, tert-butyl ester.

To a solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-4-iodo-(L)-phenylalanine tert-butyl ester (100 mg, 0.15 mmol) in toluene (1 mL) and ethanol (0.5 mL) were added 4-fluorobenzeneboronic acid (24 mg, 0.16 mmol), potassium bromide (20 mg, 0.17 mmol), 2M  $\text{Na}_2\text{CO}_3$  (0.20 mL, 0.38 mmol), and tetrakis(triphenylphosphine)palladium (9 mg, 0.008 mmol). The mixture was stirred for 1.5 hours at  $95^\circ\text{C}$  under a nitrogen atmosphere, allowed to cool to room temperature, diluted with ethyl acetate, washed twice with 1N sodium hydroxide, once with saturated brine solution, dried ( $\text{MgSO}_4$ ), and evaporated. The title compound was obtained pure by silica gel chromatography eluting with 10% acetone in hexane; yield 36 mg (38%).

MS: m/e 621 (M + H); 638 (M + H +  $\text{NH}_3$ )  
20 400 MHz  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.47 (s, 9H); 1.65-1.87 (m, 4H); 3.08 (dd, 1H); 3.20 (dd, 1H); 3.45 (m, 1H); 4.24 (dd, 1H); 4.63 (dd, 1H); 7.15 (t, 2H); 7.35 (d, 2H); 7.54 (d, 2H); 7.57 (m, 2H); 7.77-7.80 (m, 3H).

25 Step E: N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4'-fluorophenyl)phenylalanine.

A cooled solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-4-(4-fluorophenyl)-(L)-phenylalanine tert-butyl ester (36 mg, 0.055 mmol) in methylene chloride (1.4 mL) was treated with trifluoroacetic acid (0.28 mL, 3.63 mmol). The cooling bath was removed, and the mixture was stirred overnight at room temperature. The reaction mixture was then evaporated, coevaporated with methylene chloride (3X), toluene (2X), and finally methanol. The product was dried under high vacuum; yield 32 mg.

MS: m/e 565 (M + H); 582 (M + H + NH<sub>3</sub>)  
400 MHz <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 1.60-1.90 (m, 4H); 3.10 (dd, 1H); 3.42 (m, 1H); 4.22 (t, 1H); 4.73 (m, 1H); 7.11 (t, 2H); 7.34 (d, 2H); 7.52 (d, 2H); 7.56 (m, 2H); 7.72-7.79 (m, 3H).

5

## EXAMPLE 4

10 N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-thienyl)-phenylalanine

This compound was prepared in a similar fashion as Example 3 using 2-thienyl-boronic acid in the Suzuki coupling reaction; MS: m/e 553 (M + 1); 570 (M + 1 + NH<sub>3</sub>).

15

## EXAMPLE 5

20 N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-thienyl)-phenylalanine

This compound was prepared in a similar fashion as Example 3 using 3-thienyl-boronic acid in the Suzuki coupling reaction; MS: m/e 553 (M + 1); 570 (M + 1 + NH<sub>3</sub>).

25

## EXAMPLE 6

30 N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4'-trifluoromethyl-phenyl)-phenylalanine

This compound was prepared in a similar fashion as Example 3 using 4-trifluoromethylbenzene boronic acid in the Suzuki coupling reaction; MS: m/e 615 (M + 1); 632 (M + 1 + NH<sub>3</sub>).

## EXAMPLE 7

N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methoxy-phenyl)-phenylalanine

5

This compound was prepared in a similar fashion as Example 3 using 2-methoxybenzene boronic acid in the Suzuki coupling reaction; MS: m/e 577 (M + 1); 594 (M + 1 + NH<sub>3</sub>).

10

## EXAMPLE 8

N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-formyl-phenyl)-phenylalanine

15

This compound was prepared in a similar fashion as Example 3 using 2-formyl-benzene boronic acid in the Suzuki coupling reaction; MS: m/e 575 (M + 1); 592 (M + 1 + NH<sub>3</sub>).

20

The following compounds were also prepared by the procedures described in Example 3 using the appropriate boronic acid derivative in Step D:

Ex. No.	Name	Mass Spectrum
(9)	N-(3-fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-thienyl)phenylalanine;	503 (M + 1); 520 (M+1+ NH <sub>3</sub> )
(10)	N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2',6'-difluorophenyl)phenylalanine;	600 (M+1+ NH <sub>3</sub> )
(11)	N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-hydroxymethylphenyl) phenylalanine;	594 (M+1+ NH <sub>3</sub> )

- |  |  |
|--|--|
| (12) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4'-methylphenyl)phenylalanine;          | 561 (M + 1);<br>578 (M+1+NH <sub>3</sub> ) |
| (13) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-carboxyphenyl)phenylalanine;         | 608 (M+1+NH <sub>3</sub> )                 |
| (14) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methoxycarbonylphenyl)phenylalanine; | 605 (M + 1);<br>622 (M+1+NH <sub>3</sub> ) |
| (15) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-formylphenyl)phenylalanine;          | 575 (M + 1);<br>592 (M+1+NH <sub>3</sub> ) |
| (16) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-aminophenyl)phenylalanine;           | 562 (M + 1);<br>579 (M+1+NH <sub>3</sub> ) |
| (17) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylphenyl)phenylalanine;          | 561 (M + 1);<br>578 (M+1+NH <sub>3</sub> ) |
| (18) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-acetamidophenyl)phenylalanine;       | 621 (M+1+NH <sub>3</sub> )                 |
| (19) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-fluorophenyl)phenylalanine;          | 565 (M + 1);<br>582 (M+1+NH <sub>3</sub> ) |
| (20) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-carboxyphenyl)phenylalanine;         | 608 (M+1+NH <sub>3</sub> )                 |
| (21) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-methoxycarbonylphenyl)phenylalanine; | 622 (M+1+NH <sub>3</sub> )                 |
| (22) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2',4'-dichlorophenyl)phenylalanine;     | 632 (M + NH <sub>4</sub> )                 |

- (23) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl)-(L)-4-(2'-formyl-3-thienyl)phenylalanine; 581 (M + 1)

The following compounds were prepared by the procedures described in Example 3 using a 2(S)-methylproline derivative in Step B and the appropriate boronic acid derivative in Step D:

5

Ex. No.	Name	Mass Spectrum
(24)	N-(3,5-dichlorobenzenesulfonyl)-2(S)-methylprolyl-(L)-4-(4'-fluorophenyl)phenylalanine;	596 (M + NH <sub>3</sub> )
(25)	N-(3,5-dichlorobenzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-formylphenyl)phenylalanine;	606 (M + NH <sub>3</sub> )
(26)	N-(3,5-dichlorobenzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-(hydroxymethyl)phenyl)phenylalanine;	608 (M + NH <sub>3</sub> )
(27)	N-(3,5-dichlorobenzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-cyanophenyl)phenylalanine;	586 (M + 1); 603 (M + NH <sub>4</sub> )
(28)	N-(benzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-formylphenyl)phenylalanine;	521 (M + 1)
(29)	N-(benzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-methoxyphenyl)phenylalanine;	523 (M + 1)

### EXAMPLE 30

N-(3-Fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine

5 Step A: N-(3-Fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-trimethylstannylphenylalanine, tert-butyl ester.

A solution of N-(3-fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-iodophenylalanine, tert-butyl ester (prepared according to the method described in Example 3) (1.0 gm, 1.53 mmol),  
10 hexamethylditin (411  $\mu$ L, 2.14 mmol), triphenylphosphine (8 mg, 0.03 mmol), lithium chloride (71 mg, 1.68 mmol), and tetrakis(triphenylphosphine)palladium(0) (88 mg, 0.077 mmol) in 1,4-dioxane (10 mL) was heated to 95°C under a dry nitrogen atmosphere for 1.5 hr. The solution was cooled and diluted with  
15 ethyl acetate (100 mL) and successively washed with 1N sodium hydroxide solution (2X) and saturated salt solution (1X). After drying over anhydrous magnesium sulfate, the solution was filtered and the solvent removed by rotoevaporation. The residue was  
20 purified by silica gel column chromatography eluted with 10% acetone in hexanes to yield N-(3-fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(trimethylstannyl)phenylalanine, tert-butyl ester (577 mg, 54% yield).  
MS: m/e 658 (M + 18;  $\text{NH}_4^+$ ).

25 Step B: N-(3-Fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine, tert-butyl ester.

To a solution of N-(3-fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(trimethylstannyl)phenylalanine, tert-butyl ester (50 mg, 0.078 mmol) in toluene (2 mL) was added 2-bromobenzonitrile (14 mg, 0.078  
30 mmol). The solution was degassed under a dry nitrogen atmosphere (3X). Dichlorobis(triphenylphosphine)palladium(II) (2 mg, 0.0023 mmol) was added and the reaction heated to 100°C for 2 hr. Additional 2-bromobenzonitrile (7 mg, 0.039 mmol) and dichlorobis(triphenylphosphine)palladium(II) (2 mg, 0.0023 mmol) was

added and the reaction continued to be heated for 1 hr. The reaction was cooled and ethyl acetate added. The solution was washed with water and saturated salt solution and dried over anhydrous magnesium sulfate. The solvent was removed by rotoevaporation and the residue purified by silica gel column chromatography eluted with 20% acetone in hexanes to yield N-(3-fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine, tert-butyl ester (24 mg, 53% yield).

Step C: N-(3-Fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine

To a solution of N-(3-fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine, tert-butyl ester (24 mg, 0.042 mmol) in ice cooled methylene chloride (1 mL) was added trifluoroacetic acid (198  $\mu$ L, 2.58 mmol). The ice bath was removed and the solution stirred at room temperature overnight. The solvent was removed by rotoevaporation and then coevaporated with methylene chloride (2X), toluene (2X) and methanol (2X) to yield N-(3-fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine as a solid (21.5 mg, 98% yield).

MS: m/e 522 (M + 1), 539 (M + 18;  $\text{NH}_4^+$ ).

400 MHz  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.51-1.87 (m, 4H); 3.12-3.26 (m, 2H); 4.17 (dd, 1H); 4.78 (m, 1H); 7.41-7.81 (m, 12H); 8.17 (d, 1H).

The following compounds were also prepared by analogous procedures described in Example 28 using the appropriate acylating or sulfonylating agent in step A and the appropriate aryl halide derivative in Step B:

Ex. No.	Name	Mass Spectrum
(31)	N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4'-fluoro-2'-methoxyphenyl)phenylalanine;	612 (M + $\text{NH}_4$ )

- |  |                            |
|--|----------------------------|
| (32) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;   | 589 (M + NH <sub>4</sub> ) |
| (33) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylthio-phenyl)phenylalanine;   | 610 (M + NH <sub>4</sub> ) |
| (34) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(5-methyl-1,3,4-oxadiazol-2-yl-phenyl)phenyl)phenylalanine;              | 629 (M + 1)                |
| (35) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-methyl-5-trifluoromethyl-benzoxazol-7-yl)phenylalanine;                   | 670 (M + 1)                |
| (36) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-methyl-6-(5-trifluoromethyl-tetrazol-1-yl)-benzoxazol-4-yl)phenylalanine; | 738 (M + 1)                |
| (37) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-methyl-5-(5-trifluoromethyl-tetrazol-1-yl)-benzoxazol-7-yl)phenylalanine; | * see NMR below            |
| (38) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3-pyridyl)phenylalanine;  | 548 (M + 1)                |
| (39) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-pyridyl)phenylalanine;  | 548 (M + 1)                |
| (40) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(5-pyrimidinyl)phenylalanine;  | 549 (M + 1)                |
| (41) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-cyanophenyl)phenylalanine;   | 589 (M + NH <sub>4</sub> ) |
| (42) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-methyl-benzoxazol-4-yl)phenylalanine;                                     | 602 (M + 1)                |



- (43) N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(6-acetamido-2-methylbenzoxazol-4-yl)phenylalanine; 659 (M + 1)
- (44) N-(benzenesulfonyl)-(L)-prolyl-(L)-4-(2-pyridyl)phenylalanine;
- (45) N-(3,5-dichlorobenzenesulfonyl)-(L)-3(S)-methylprolyl-(L)-4-(2'-cyanophenyl)phenylalanine; 603 (M + NH<sub>4</sub>)

\* Example 37: 400 MHz <sup>1</sup>H NMR (CD<sub>3</sub>OD): d 1.6-1.95(m, 4H), 2.70(s, 3H), 3.1-3.2(dd, 1H), 3.3-3.45(m, 3H), 4.2-4.3(m, 1H), 4.72-4.8(m, 1H), 7.48 (d, 2H), 7.68-7.9 (m, 6H), 8.3(d, 1H)

5

#### EXAMPLE 46

N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(tetrazol-5-yl)phenyl)phenylalanine.

- 10 Step A: N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(tetrazol-5-yl)phenyl)phenylalanine, tert-butyl ester.  
Trimethyltinazide (115 mg, 0.556 mmol) was added to a solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine, tert-butyl ester (prepared according to the procedures described in Example 27, Step B) (100 mg, 0.159 mmol) in toluene under dry nitrogen atmosphere. The solution was heated to 115° C for 18 hr. Upon cooling to room temperature, the solvent was removed by rotoevaporation and the residue dissolved in ethyl acetate. The solution was successively washed with 5N hydrochloric acid and saturated salt solution and dried over anhydrous sodium sulfate. The mixture was filtered and the solvent removed by rotoevaporation. The residue was purified by flash column chromatography on silica gel eluted with 2-5% methanol in methylene chloride to yield N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(tetrazol-5-yl)phenyl)phenylalanine, tert-butyl ester (32 mg, 30% yield). MS: m/e 671 (M + 1).

Step B: N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(tetrazol-5-yl)phenyl)phenylalanine.

- Trifluoroacetic acid (0.227 mL, 2.95 mmol) was added to a solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(tetrazol-5-yl)phenyl)phenylalanine, tert-butyl ester (32 mg, 0.048 mmol) in methylene chloride (1.2 mL) and stirred for 18 hr at room temperature. The solution was concentrated by rotoevaporation to a solid and then successively co-evaporated with methylene chloride, toluene and methanol. The crude solid was purified by flash column chromatography on silica gel eluted with 0.5% acetic acid in 5% methanol / methylene chloride to give N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(tetrazol-5-yl)phenyl)phenylalanine (15mg).
- MS: m/e 615 (M + 1).  
400 MHz <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 3.44 (m, 1H), 4.19 (t, 1H) 4.68 (t, 1H), 7.07 (d, 2H), 7.23 (d, 2H), 7.52 (m, 2H), 7.64 (m, 2H), 7.78 (m, 3H).

20

#### EXAMPLE 47

N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(2-methyl-tetrazol-5-yl)phenyl)phenylalanine.

- Step A. N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(2-methyl-tetrazol-5-yl)phenyl)phenylalanine, tert-butyl ester  
Potassium carbonate (10 mg, 0.075 mmol) and methyl iodide (4.6 μL, 0.075 mmol) were added to a solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(tetrazol-5-yl)phenyl)phenylalanine, tert-butyl ester (from Example 36, Step A) (33 mg, 0.049 mmol) in dimethylformamide and stirred at room temperature for 2 hr. The reaction was partitioned between water and ethyl acetate and separated. The organic layer was successively washed with water and saturated salt solution and dried over anhydrous sodium

sulfate. The solvent was removed by rotoevaporation and the crude solid purified by flash column chromatography on silica gel eluted with 10% acetone in hexanes. The component that eluted first proved to be N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(1-methyl-tetrazol-5-yl)phenyl)phenylalanine, tert-butyl ester (8 mg, 24% yield).  
MS: m/e 685 (M + 1).

The component that eluted second was N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(2-methyl-tetrazol-5-yl)phenyl)phenylalanine, tert-butyl ester (17 mg, 51% yield).

MS: m/e 685 (M + 1).

Step B. N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(2-methyl-tetrazol-5-yl)phenyl)phenylalanine.

Trifluoroacetic acid (56  $\mu$ L, 0.72 mmol) was added to a solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(2-methyl-tetrazol-5-yl)phenyl)phenylalanine, tert-butyl ester (8 mg, 0.012 mmol) in methylene chloride (0.5 mL) and stirred for 18 hr at room temperature. The solvent was removed by rotoevaporation and the crude solid successively co-evaporated with methylene chloride, toluene, and methanol to yield N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(2-methyl-tetrazol-5-yl)phenyl)phenylalanine as a white solid (6.5 mg, 86% yield).

MS: m/e 629 (M + 1).

400 MHz  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  3.45 (m, 1H), 4.23 (t, 1H), 4.27 (s, 3H), 7.07 (d, 2H), 7.19 (d, 2H), 7.46 (m, 2H), 7.70 (d, 1H), 7.78 (d, 3H), 8.14 (d, 1H).

#### EXAMPLE 48

N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(3-methyl-tetrazol-5-yl)phenyl)phenylalanine.

Trifluoroacetic acid (120  $\mu$ L, 1.54 mmol) was added to a solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(3-

5 methyl-tetrazol-5-yl)phenyl)phenylalanine, tert-butyl ester (from Example 37, Step A) (17 mg, 0.025 mmol) in methylene chloride (0.6 mL) and stirred for 18 hr at room temperature. The solvent was removed by rotoevaporation and the crude solid successively co-evaporated with methylene chloride, toluene, and methanol. The crude solid was purified by flash column chromatography on silica gel eluted with 0-0.5% acetic acid in 5% methanol / methylene chloride to yield N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(2-methyl-tetrazol-5-yl)phenyl)phenylalanine as a white solid (9 mg, 57% yield).

10 MS: m/e 629 (M + 1).  
400 MHz <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 3.16 (m, 1H), 3.30 (s, 3H), 4.23 (m, 1H), 4.65 (m, 1H), 7.05 (d, 2H), 7.27 (d, 2H), 7.60 (m, 3H), 7.72 (m, 1H), 7.80 (d, 3H), 8.17 (d, 1H).

#### 15 EXAMPLE 49

N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-aminocarbonylphenyl)phenylalanine.

20

Step A: N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-carboxyphenyl)phenylalanine.

A solution of tetrabutylammonium permanganate (465 mg, 1.28 mmol) in pyridine (8 mL) was added to a solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-formylphenyl)phenylalanine, tert-butyl ester (from Example 8) (810 mg, 1.28 mmol) in pyridine. The deep purple solution was stirred for 1.5 hr at room temperature and then poured into an ice cold solution of sodium sulfite (7.5 gm) in 5N hydrochloric acid to yield a white precipitate. The mixture was extracted with ethyl acetate (3 x 50 mL). The organic extracts were dried over anhydrous sodium sulfate, filtered and the solvent removed by rotoevaporation to yield a white solid. This solid was purified by flash column chromatography on silica gel eluted with 5% methanol in methylene chloride to yield N-(3,5-

25  
30

dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-carboxyphenyl)phenylalanine, tert-butyl ester (550 mg, 66% yield).  
400 MHz  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.45 (s, 9H), 3.47 (m, 1H), 4.20 (m, 1H), 4.63 (m, 1H), 7.29 (m, 5H), 7.39 (t, 1H), 7.50 (t, 1H), 7.74 (d, 1H), 7.80 (m, 3H), 8.21 (d, 1H).

Step B: N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-aminocarbonylphenyl)phenylalanine, tert-butyl ester.

To a solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-carboxyphenyl)phenylalanine, tert-butyl ester (400 mg, 0.618 mmol) in tetrahydrofuran (4 mL) cooled to  $-5^\circ\text{C}$  was added N-methylmorpholine (68  $\mu\text{L}$ , 0.618 mmol) and isobutyl chloroformate (80  $\mu\text{L}$ , 0.618 mmol). The solution was stirred for 5 min and then an aqueous solution of 30% ammonium hydroxide (0.10 mL, 0.928 mmol) was added. After stirring for 1 hr at room temperature, the mixture was concentrated by rotoevaporation. The residue was purified by flash column chromatography on silica gel eluted with 2-5% methanol in methylene chloride to yield N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-aminocarbonylphenyl)phenylalanine, tert-butyl ester (110 mg).

Step C: N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-aminocarbonylphenyl)phenylalanine.

Trifluoroacetic acid (148  $\mu\text{L}$ , 1.92 mmol) was added to a solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-aminocarbonylphenyl)phenylalanine, tert-butyl ester (110 mg, 0.17 mmol) and stirred for 18 hr at room temperature. The reaction was concentrated by rotoevaporation and the solid successively co-evaporated with methylene chloride, toluene and methanol. The residue was purified by flash column chromatography on silica gel eluted with 0-0.5% acetic acid in 5% methanol / methylene chloride to give N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-aminocarbonylphenyl)phenylalanine as a white solid (14 mg).  
MS: m/e 590 (M + 1).

400 MHz  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  3.45 (m, 1H), 4.22 (m, 1H), 4.71 (t, 1H), 7.30-7.40 (m, 6H), 7.46 (t, 1H), 7.53 (d, 1H), 7.78 (d, 3H).

The following compounds were also prepared by  
5 analogous procedures described in Example 39:

Ex. No.	Name	MS
(50)	N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylaminocarbonylphenyl)phenylalanine;	604 (M + 1); 621 (M + $\text{NH}_4$ )
(51)	N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;	618 (M + 1), 635 (M + $\text{NH}_4$ )
(52)	N-(benzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-carboxyphenyl)phenylalanine;	554 (M + $\text{NH}_4$ )

### EXAMPLE 53

10

N-(1-Butanesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine

Step A : N-(tert-Butyloxycarbonyl)-(L)-prolyl-(L)-4-iodophenylalanine, tert-butyl ester

15

To a solution of N-(tert-butyloxycarbonyl)-(L)-proline (1.12 g, 5.20 mmol) in methylene chloride (100 mL) were added 1-hydroxybenzotriazole (1.04 g, 7.70 mmol), N-methylmorpholine (1.4 mL, 12.7 mmol), and 4-iodo-(L)-phenylalanine, tert-butyl ester hydrochloride (2.0 g, 5.21 mmol). After cooling in an ice-bath for 5 minutes, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (1.19 g, 6.21 mmol) was added. After 15 minutes, the cooling bath was removed, and the mixture was stirred overnight under a nitrogen atmosphere. The mixture was diluted with methylene chloride, successively washed with water, 1N hydrochloric acid, saturated sodium

20

bicarbonate solution, and saturated salt solution, and dried over anhydrous magnesium sulfate. The solvent was removed by rotoevaporation and the residue purified by flash column chromatography on silica gel eluted with 25% ethyl acetate in hexane to afford N-(*tert*-butyloxycarbonyl)-(L)-prolyl-(L)-4-iodophenylalanine, *tert*-butyl ester (2.37 gm, 84% yield).

Step B: N-(*tert*-Butyloxycarbonyl)-(L)-prolyl-(L)-4-(trimethylstannyl)phenylalanine, *tert*-butyl ester

This intermediate was prepared following the procedure described in Step A of Example 28, using N-(*tert*-butyloxycarbonyl)-(L)-prolyl-(L)-4-iodophenylalanine, *tert*-butyl ester as starting material. The compound was obtained pure by flash column chromatography on silica gel eluted with 20% ethyl acetate in hexane (69% yield).

Step C: N-(*tert*-Butyloxycarbonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine, *tert*-butyl ester

This intermediate was prepared following the procedure described in Step B of Example 28, using N-(*tert*-butyloxycarbonyl)-(L)-prolyl-(L)-4-(trimethylstannyl)phenylalanine, *tert*-butyl ester as starting material and purified by flash column chromatography in silica gel eluted with 20% ethyl acetate in hexane.

Step D: (L)-Prolyl-(L)-4-(2'-cyanophenyl)phenylalanine, *tert*-butyl ester hydrochloride

N-(*tert*-Butyloxycarbonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine, *tert*-butyl ester (1.11 g, 2.14 mmol) was stirred with 1M hydrochloric acid in ethyl acetate (10.6 mL) overnight at room temperature. The reaction mixture was rotoevaporated and co-evaporated several times with diethyl ether. Flash column chromatography on silica gel eluted with 5% methanol in methylene

chloride afforded (L)-Prolyl-(L)-4-(2'-cyanophenyl)phenylalanine, *tert*-butyl ester hydrochloride (742 mg, 76% yield).

400 MHz  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.44 (s, 9H); 3.08 (dd, 1H); 3.18-3.31 (m, 3H); 4.11 (dd, 1H); 4.67 (dd, 1H); 7.39-7.82 (m, 8H).

5

Step E: N-(1-Butanesulfonyl)-(L-)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine, *tert*-butyl ester

To a solution of (L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine, *tert*-butyl ester hydrochloride (45 mg, 0.099 mmol) in methylene chloride (2 mL) were added N,N-diisopropylethylamine (52  $\mu\text{L}$ , 0.299 mmol), 4-dimethylaminopyridine (2 mg, 0.016 mmol), and 1-butan sulfonyl chloride (20  $\mu\text{L}$ , 0.154 mmol). The reaction mixture was stirred overnight at room temperature, diluted with methylene chloride, successively washed with water, 2N hydrochloric acid, saturated sodium bicarbonate solution, and saturated salt solution. After drying over anhydrous magnesium sulfate, the solvent was removed by rotoevaporation and the residue purified by flash column chromatography on silica gel eluted with 15-20% acetone in hexane to afford N-(1-butan sulfonyl)-(L-)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine, *tert*-butyl ester (24.4 mg, 46% yield). 400 MHz  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  0.92 (t, 3H); 1.45 (s, 9H); 3.01 (dd, 1H); 3.25 (dd, 1H); 4.29 (dd, 1H); 4.67 (dd, 1H); 7.38-7.82 (m, 8H).

25

Step F: N-(1-Butanesulfonyl)-(L-)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine

A cooled solution of N-(1-butan sulfonyl)-(L-)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine, *tert*-butyl ester (22 mg, 0.041 mmol) in methylene chloride (2.0 mL) was treated with trifluoroacetic acid (0.10 mL, 1.30 mmol). The cooling bath was removed, and the mixture was stirred overnight at room temperature. The reaction mixture was then rotoevaporated, co-evaporated with methylene chloride (3X), toluene



(2X), and finally methanol. The residue was purified by flash column chromatography on silica gel eluted with 0-0.1% acetic acid in 3% methanol in methylene chloride to afford N-(1-butanesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine (17 mg).

- 5 MS: m/e 484 (M + H); 501 (M + NH<sub>4</sub>);  
400 MHz <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 0.90 (t, 3H); 1.40 (m, 2H); 3.03 (m, 2H); 3.13 (dd, 1H); 4.29 (dd, 1H); 4.77 (m, 1H); 7.38-7.82 (m, 8H).

- 10 The following compounds were also prepared by analogous procedures described in Example 39 using the appropriate acyl or sulfonyl halide derivative in Step B:

Ex. No.	Name	MS
(54)	N-(3-bromobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;	601 (M + NH <sub>4</sub> )
(55)	N-(benzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;	504 (M + 1); 521 (M + NH <sub>4</sub> )
(56)	N-(α-toluenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;	518 (M + 1); 535 (M + NH <sub>4</sub> )
(57)	N-(phenylacetyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;;	482 (M + 1); 499 (M + NH <sub>4</sub> )
(58)	N-(3-pyridinesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;	505 (M + 1)
(59)	N-(2-thienylsulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;	527 (M + NH <sub>4</sub> )
(60)	N-(benzylaminocarbonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;	497 (M + 1)
(61)	N-(3-phenylpropionyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;	496 (M + 1)
(62)	N-((5-methyl-3,4-thiadiazol-2-yl)sulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;	543 (M + NH <sub>4</sub> )

- (63) N-((benzthiazol-2-yl)sulfonyl)-(L)-prolyl- 578 (M + NH<sub>4</sub>)  
(L)-4-(2'-cyanophenyl)phenylalanine;  
(64) N-((1-methyl-imidazol-4-yl)sulfonyl)-(L)- 508 (M + 1)  
prolyl-(L)-4-(2'-  
cyanophenyl)phenylalanine;

### EXAMPLE 65

5 N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-  
methylsulfinylphenyl)phenylalanine.

Step A: N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-  
methylsulfinylphenyl)phenylalanine, tert-butyl ester.

10 To a solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-  
prolyl-(L)-4-(2'-methylthiophenyl)phenylalanine, tert-butyl ester  
(from Example 33 and 28) (22 mg, 0.034 mmol) in methylene  
chloride ( 1.0 mL) was added 3-chloroperoxybenzoic acid (12 mg,  
0.034 mmol). After stirring the solution at room temperature for 15  
15 min, solid sodium bisulfite (25 mg) was added and the solvent  
removed by rotoevaporation. The remaining solid was purified by  
flash column chromatography on silica gel eluted with 25% acetone  
in hexane to afford N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-  
4-(2'-methylsulfinylphenyl)phenylalanine, tert-butyl ester (15 mg)  
20 which was used in the subsequent reaction.

Step B: N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-  
(2'-methylsulfinylphenyl)phenylalanine.

25 Trifluoroacetic acid (111 µL) was added to an ice cooled  
solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-  
methylsulfinylphenyl)phenylalanine, tert-butyl ester (15 mg) in  
methylene chloride (1.0 mL). After stirring for 18 hr at room  
temperature, the solvent was removed by rotoevaporation to yield N-  
(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-

methylsulfinylphenyl)phenylalanine (14.6 mg) as a 1:1 mixture of sulfoxide diastereomers.

MS:  $m/e = 609$  ( $M + 1$ ).

400 MHz  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  2.43, 2.44 (s, 3H), 3.46 (m, 1H),

5 4.23 (m, 1H), 4.74 (m, 1H), 7.35 (m, 3H), 7.42 (m, 2H), 7.61 (m, 3H), 7.79 (s, 2H), 8.00 (d, 1H), 8.32 (d, 1H).

### EXAMPLE 66

10

N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylsulfonylphenyl)phenylalanine.

15

Step A: N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylsulfonylphenyl)phenylalanine, tert-butyl ester.

20

To a solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylthiophenyl)phenylalanine, tert-butyl ester (from Example 33 and 28) (22 mg, 0.034 mmol) in methylene chloride (1.0 mL) was added 3-chloroperoxybenzoic acid (24 mg, 0.068 mmol) in two equal portions 15 min apart. After stirring the solution at room temperature for 4 hr, solid sodium bisulfite (25 mg) was added and the solvent removed by rotoevaporation. The remaining solid was purified by flash column chromatography on silica gel eluted with 25% acetone in hexane to afford N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylsulfonylphenyl)phenylalanine, tert-butyl ester (15 mg) which was used in the subsequent reaction.

25

30

Step B: N-(3,5-Dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylsulfonylphenyl)phenylalanine.

Trifluoroacetic acid (111  $\mu\text{L}$ ) was added to an ice cooled solution of N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylsulfonylphenyl)phenylalanine, tert-butyl ester (15 mg) in methylene chloride (1.0 mL). After stirring for 18 hr at room

temperature, the solvent was removed by rotoevaporation to yield N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylsulfonylphenyl)phenylalanine (14.3 mg).

MS:  $m/e = 624$  (M), 642 (M + NH<sub>4</sub>).

5 400 MHz <sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  2.65, 2.44 (s, 3H), 3.50 (m, 1H), 4.22 (m, 1H), 4.75 (m, 1H), 7.39 (m, 3H), 7.48 (t, 1H), 7.61 (m, 1H), 7.69 (t, 1H), 7.81 (d, 3H), 7.97 (m, 1H), 8.15 (d, 1H), 8.28 (d, 1H).

10

### EXAMPLE 67

#### Inhibition of VLA-4 Dependent Adhesion to BSA-CS-1 Conjugate

##### Step 1. Preparation of CS-1 Coated Plates

15

Untreated 96 well polystyrene flat bottom plates were coated with bovine serum albumin (BSA; 20  $\mu$ g/ml) for 2 hours at room temperature and washed twice with phosphate buffered saline (PBS).

20 The albumin coating was next derivatized with 10  $\mu$ g/ml 3-(2-pyridyldithio) propionic acid N-hydroxysuccinimide ester (SPDP), a heterobifunctional crosslinker, for 30 minutes at room temperature and washed twice with PBS. The CS-1 peptide (Cys-Leu-His-Gly-Pro-Glu-Ile-Leu-Asp-Val-Pro-Ser-Thr), which was synthesized by conventional solid phase chemistry and purified by reverse phase HPLC, was next  
25 added to the derivatized BSA at a concentration of 2.5  $\mu$ g/ml and allowed to react for 2 hours at room temperature. The plates were washed twice with PBS and stored at 4°C.

##### Step 2. Preparation of Fluorescently Labeled Jurkat Cells

30

Jurkat cells, clone E6-1, obtained from the American Type Culture Collection (Rockville, MD; cat # ATCC TIB-152) were grown and maintained in RPMI-1640 culture medium containing 10% fetal calf serum (FCS), 50 units/ml penicillin, 50  $\mu$ g/ml streptomycin and 2 mM

L-Glutamine. Fluorescence activated cell sorter analysis with specific monoclonal antibodies confirmed that the cells expressed both the  $\alpha 4$  and  $\beta 1$  chains of VLA-4. The cells were centrifuged at 400xg for five minutes and washed twice with PBS. The cells were incubated at a  
5 concentration of  $2 \times 10^6$  cells/ml in PBS containing a 1  $\mu$ M concentration of a fluorogenic esterase substrate (2', 7'-bis-(2-carboxyethyl)-5-(and -6)-carboxyfluorescein, acetoxymethyl ester; BCECF-AM; Molecular Probes Inc., Eugene, Oregon; catalog #B-1150) for 30-60 minutes at 37°C in a 5% CO<sub>2</sub>/air incubator. The  
10 fluorescently labeled Jurkat cells were washed two times in PBS and resuspended in RPMI containing 0.25% BSA at a final concentration of  $2.0 \times 10^6$  cells/ml.

Step 3. Assay Procedure

15

Compounds of this invention were prepared in DMSO at 100x the desired final assay concentration. Final concentrations were selected from a range between 0.001 nM-100mM. Three mL of diluted compound, or vehicle alone, were premixed with 300 mL of cell  
20 suspension in 96-well polystyrene plates with round bottom wells. 100 mL aliquots of the cell /compound mixture were then transferred in duplicate to CS-1 coated wells. The cells were next incubated for 30 minutes at room temperature. The non-adherent cells were removed by two gentle washings with PBS. The remaining adherent cells were  
25 quantitated by reading the plates on a Cytofluor II fluorescence plate reader (Perseptive Biosystems Inc., Framingham, MA; excitation and emission filter settings were 485 nm and 530 nm, respectively). Control wells containing vehicle alone were used to determine the level of cell adhesion corresponding to 0% inhibition. Wells in which cells were  
30 treated with a saturating concentration (10 ng/ml) of a neutralizing anti- $\alpha 4$  antibody (HP 2/1; Immunotech, Inc., Westbrook, ME) were used to determine the level of cell adhesion corresponding to 100% inhibition. Cell adhesion in the presence of HP2/1 was usually less than 5% of that observed in the presence of vehicle alone. Percent inhibition was then

calculated for each test well and the IC<sub>50</sub> was determined from an eight point titration using a validated four parameter fit algorithm. IC<sub>50</sub> values for the inhibition of VCAM-Ig binding to VLA-4 are provided below for representative compounds:

- 5 <100 nM - compounds of examples 15, 20, 37, 62;  
<10 nM - compound of example 27, 32, 46, 55, 58, 65, 66.

### EXAMPLE 68

10

#### Antagonism of VLA-4 Dependent Binding to VCAM-Ig Fusion Protein.

##### 1. Preparation of VCAM-Ig

15

The signal peptide as well as domains 1 and 2 of human VCAM (GenBank Accession no. M30257) were amplified by PCR using the human VCAM cDNA (R & D Systems) as template and the following primer sequences: 3'-PCR primer:5'-

AATTATAATTTGATCAACTTAC

20

CTGTCAATTCTTTTACAGCCTGCC-3';

5'-PCR primer:

5'-ATAGGAATTCCAGCTGCCACCATGCCTGGGAAGATGGTTCG-3'.

25 The 5'-PCR primer contained EcoRI and PvuII restriction sites followed by a Kozak consensus sequence (CCACC) proximal to the initiator methionine ATG. The 3'-PCR primer contained a BclI site and a splice donor sequence. PCR was performed for 30 cycles using the following parameters: 1 min. at 94°C, 2 min. at 55°C and 2 min. at 72°C. The amplified region encoded the following sequence of human VCAM:

30

MPGKMVVILGASNILWIMFAASQAFKIETTPESRYLAQIGDSVSLT  
CSTTGCESPFFSWRTQIDSPLNGKVTNEGTTSTLTMNPVSFGNEHS  
YLCTATCESRKLEKGIQVEIYSFPKDPEIHLSGPLEAGKPITVKCSV  
ADVYPFDRLEIDLLKGDHLMKSQEFLEDADRKSLETKSLEVTFTP  
VIEDIGKVLVCRAKLHIDEMDSVPTVRQAVKEL. The resulting

PCR product of 650 bp was digested with EcoRI and BclI and ligated to expression vector pIg-Tail (R & D Systems, Minneapolis, MN) digested with EcoRI and BamHI. The pIg-Tail vector contains the genomic fragment which encodes the hinge region, CH2 and CH3 of human IgG1 (GenBank Accession no. Z17370). The DNA sequence of the resulting VCAM fragment was verified using Sequenase (US Biochemical, Cleveland, OH). The fragment encoding the entire VCAM-Ig fusion was subsequently excised from pIg-Tail with EcoRI and NotI and ligated to pCI-neo (Promega, Madison, WI) digested with EcoRI and NotI. The resulting vector, designated pCI-neo/VCAM-Ig was transfected into CHO-K1 (ATCC CCL 61) cells using calcium-phosphate DNA precipitation (Specialty Media, Lavalette, NJ). Stable VCAM-Ig producing clones were selected according to standard protocols using 0.2-0.8 mg/ml active G418 (Gibco, Grand Island, NY), expanded, and cell supernatants were screened for their ability to mediate Jurkat adhesion to wells previously coated with 1.5 µg/ml (total protein) goat anti-human IgG (Sigma, St. Louis, MO). A positive CHO-K1/VCAM-Ig clone was subsequently adapted to CHO-SFM serum-free media (Gibco) and maintained under selection for stable expression of VCAM-Ig. VCAM-Ig was purified from crude culture supernatants by affinity chromatography on Protein A/G Sepharose (Pierce, Rockford, IL) according to the manufacturer's instructions and desalted into 50 mM sodium phosphate buffer, pH 7.6, by ultrafiltration on a YM-30 membranes (Amicon, Beverly, MA).

25

Step 2.      Preparation of  $^{125}$ I-VCAM-Ig

VCAM-Ig was labeled to a specific radioactivity greater than 1000 Ci/mmol with  $^{125}$ I-Bolton Hunter reagent (New England Nuclear, Boston, MA; cat # NEX120-0142) according to the manufacturer's instructions. The labeled protein was separated from unincorporated isotope by means of a calibrated HPLC gel filtration column (G2000SW; 7.5 x 600 mm; Tosoh, Japan) using uv and radiometric detection.

Step 3. VCAM-Ig Binding Assay

Compounds of this invention were prepared in DMSO at  
5 100x the desired final assay concentration. Final concentrations were  
selected from a range between 0.001 nM-100 $\mu$ M. Jurkat cells were  
centrifuged at 400xg for five minutes and resuspended in binding buffer  
(25 mM HEPES, 150 mM NaCl, 3 mM KCl, 2 mM glucose, 1 mM  
MnCl<sub>2</sub>, 0.1% bovine serum albumin, pH 7.4) without MnCl<sub>2</sub>. The cells  
10 were centrifuged again and resuspended in complete binding buffer.  
Compounds were assayed in Millipore MHVB multiscreen plates (cat#  
MHVBN4550, Millipore Corp., MA) by making the following additions  
to duplicate wells: (i) 200  $\mu$ L of binding buffer; (ii) 20  $\mu$ L of a working  
stock of <sup>125</sup>I-VCAM-Ig prepared in binding buffer (final assay  
15 concentration  $\leq$  100 pM); (iii) 2.5  $\mu$ L of compound solution or vehicle  
alone; (iv) and 0.5 x 10<sup>6</sup> cells in a volume of 30  $\mu$ L. The plates were  
incubated at room temperature for 30 minutes, filtered on a vacuum  
box, and washed on the same apparatus by the addition of 100  $\mu$ L of  
binding buffer. After insertion of the multiscreen plates into adapter  
20 plates (Packard, Meriden, CT, cat# 6005178), 100  $\mu$ L of microscint-20  
(Packard cat# 6013621) was added to each well. The plates were then  
sealed, placed on a shaker for 30 seconds, and counted on a Topcount  
microplate scintillation counter (Packard). Control wells containing  
vehicle alone were used to determine the level of VCAM-Ig binding  
25 corresponding to 0% inhibition. Wells in which cells were treated with  
a saturating concentration of unlabeled VCAM-Ig (10 nM) were used to  
determine the level of binding corresponding to 100% inhibition.  
Binding of <sup>125</sup>I-VCAM-Ig in the presence of 10 nM unlabeled VCAM-  
Ig was usually less than 5% of that observed in the presence of vehicle  
30 alone. Percent inhibition was then calculated for each test well and the  
IC<sub>50</sub> was determined from a ten point titration using a validated four  
parameter fit algorithm. IC<sub>50</sub> values for the inhibition of VCAM-Ig  
binding to VLA-4 are provided below for representative compounds:  
< 100 nM: 60;



< 10 nM: 1-8, 10, 15, 20, 27, 29, 32, 37, 45, 46, 52, 55, 57, 58, 62, 64, 65, 66.

5

### EXAMPLE 69

#### Antagonism of $\alpha_4\beta_7$ Dependent Binding to VCAM-Ig Fusion Protein.

##### Step 1. $\alpha_4\beta_7$ Cell line.

10

RPMI-8866 cells (a human B cell line  $\alpha_4^+\beta_1^-\beta_7^+$ ; a gift from Prof. John Wilkins, University of Manitoba, Canada) were grown in RPMI/10% fetal calf serum/ 100 U penicillin, 100  $\mu$ g streptomycin/ 2 mM L-glutamine at 37°C, 5 % carbon dioxide. The cells ( $1.25 \times 10^6$  cells/well) were pelleted at 1000 rpm for 5 min. and then washed twice in binding buffer (25 mM Hepes, 150 mM NaCl, 0.1 % BSA, 3 mM KCl, 2 mM Glucose, pH 7.4). Cells were then resuspended at  $3.3 \times 10^7$  cells/ml.

##### Step 2. VCAM-Ig Binding Assay

20

$^{125}$ I-VCAM-Ig (prepared as above) was diluted in binding buffer to  $\leq 500$  pM VCAM-Ig /10  $\mu$ l. (i.e. if specific activity VCAM-Ig =  $3.5 \times 10^6$  cpm/pmol, then 250,000 cpm/well =  $7.14 \times 10^{-14}$  moles VCAM-Ig. The volume of the assay was 150  $\mu$ l, and the final concentration of VCAM-Ig = 476 pM). Compounds of the present invention and serial dilutions were prepared in DMSO. The final amount of DMSO in the assay was kept at 1% when adding 1.5  $\mu$ l diluted compound to each well. Compounds were assayed in Millipore MHVB multiscreen plates (Cat# MHVBN4550) by making the following sequential additions to duplicate wells: (i) 100  $\mu$ l/well of binding buffer containing 1.5 mM  $Mn^{++}$ ; (ii)  $\leq 500$  pM/well  $^{125}$ I-VCAM-Ig; (iii) 1.5  $\mu$ l/well test compound; (iv) 38  $\mu$ l/well RPMI-8866 cell suspension ( $1.25 \times 10^6$  cells/well). Control wells were established as follows: (i) total binding = buffer +  $^{125}$ I-VCAM-Ig + cells; (ii) non-specific binding =


30

buffer +  $^{125}$ I-VCAM-Ig - cells. Plates were incubated for 45 min. at 25°C on a plate shaker at 200 rpm. The multiscreen plates were filtered using a Millipore vacuum manifold (Cat. # MAVM 096 01). The plates were washed once with 100ul/well binding buffer + 1 mM  $Mn^{++}$ . After vacuum filtration, the plates were blotted, the plastic backing was removed and blotted again and allowed to air dry. When the filters were dry, the plates were transferred to Packard adapter plates (Cat# 6005178). Packard Microscint-20 (100  $\mu$ L/well) (Cat# 6013621) was added and the plates were sealed. The plates were placed on a plate shaker at 500 rpm for 30 seconds and counted on a Packard Topcount. Percent inhibition was then calculated for each test well and the IC<sub>50</sub> was determined from a ten point titration using a validated four parameter fit algorithm after subtracting the nonspecific binding values. IC<sub>50</sub> values for inhibition of VCAM-Ig binding to  $\alpha_4\beta_7$  for representative compounds are as follows:

<1000 nM: 15, 20, 27, 32, 37;  
<100 nM: 46, 55, 57, 58.

WHAT IS CLAIMED IS:

1. A compound of Formula I



5

I

or a pharmaceutically acceptable salt thereof wherein:

- 10 R<sup>1</sup> is 1) C<sub>1</sub>-10alkyl,  
 2) C<sub>2</sub>-10alkenyl,  
 3) C<sub>2</sub>-10alkynyl,  
 4) Cy,  
 5) Cy-C<sub>1</sub>-10alkyl,  
 15 6) Cy-C<sub>2</sub>-10alkenyl,  
 7) Cy-C<sub>2</sub>-10alkynyl,

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from R<sup>a</sup>; and Cy is optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

20

R<sup>2</sup> and R<sup>3</sup> are independently

- 1) hydrogen, or  
 2) a group selected from R<sup>1</sup>; or

25 R<sup>2</sup> and R<sup>3</sup> together with the atoms to which they are attached form a ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen, wherein said ring may be isolated or benzo-fused, and optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

30 R<sup>4</sup> and R<sup>7</sup> are independently selected from the group consisting of

- 1) hydrogen,  
 2) C<sub>1</sub>-10alkyl,

- 3) C<sub>2-10</sub>alkenyl,
- 4) C<sub>2-10</sub>alkynyl,
- 5) aryl,
- 6) aryl C<sub>1-10</sub>alkyl,
- 5 7) heteroaryl, and
- 8) heteroaryl C<sub>1-10</sub>alkyl,

wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents independently selected from R<sup>a</sup>, and aryl and heteroaryl are optionally substituted with one to four substituents  
 10 independently selected from R<sup>b</sup>; or

R<sub>3</sub>, R<sub>4</sub> and the carbon to which they are attached form a 3-7 membered ring optionally containing 0-2 heteroatoms selected from N, O and S;

- 15 R<sup>5</sup> is 1) hydrogen,
- 2) C<sub>1-10</sub>alkyl optionally substituted with one to four substituents independently selected from R<sup>a</sup>, or
- 3) Cy optionally substituted with one to four substituents independently selected from R<sup>b</sup>,

20

- R<sup>6</sup> is 1) Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>1-10</sub>alkyl,
- 2) Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>2-10</sub>alkenyl,
- 3) Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>2-10</sub>alkynyl,

wherein Ar<sup>1</sup> and Ar<sup>2</sup> are independently aryl or heteroaryl each  
 25 of which is optionally substituted with one to four substituents independently selected from R<sup>b</sup>; alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents independently selected from R<sup>a</sup>;

- 30 R<sup>a</sup> is 1) Cy
- 2) -OR<sup>d</sup>,
- 3) -NO<sub>2</sub>,
- 4) halogen
- 5) -S(O)<sub>m</sub>R<sup>d</sup>,

- 5           6)  $-SR^d$ ,  
              7)  $-S(O)_2OR^d$ ,  
              8)  $-S(O)_mNR^dRe$ ,  
              9)  $-NR^dRe$ ,  
              10)  $-O(CR^fRg)_nNR^dRe$ ,  
              11)  $-C(O)R^d$ ,  
              12)  $-CO_2R^d$ ,  
              13)  $-CO_2(CR^fRg)_nCONR^dRe$ ,  
              14)  $-OC(O)R^d$ ,  
 10           15)  $-CN$ ,  
              16)  $-C(O)NR^dRe$ ,  
              17)  $-NR^dC(O)Re$ ,  
              18)  $-OC(O)NR^dRe$ ,  
              19)  $-NR^dC(O)ORE$ ,  
 15           20)  $-NR^dC(O)NR^dRe$ ,  
              21)  $-CR^d(N-ORE)$ , or  
              22)  $-CF_3$ ;

wherein Cy is optionally substituted with one to four substituents independently selected from  $R^c$ ;

- 20            $R^b$  is 1) a group selected from  $R^a$ ,  
                   2)  $C_{1-10}$  alkyl,  
                   3)  $C_{2-10}$  alkenyl,  
                   4)  $C_{2-10}$  alkynyl,  
 25            5) aryl  $C_{1-10}$ alkyl,  
                   6) heteroaryl  $C_{1-10}$  alkyl,

wherein alkyl, alkenyl, alkynyl, aryl, heteroaryl are optionally substituted with a group independently selected from  $R^c$ ;

- 30            $R^c$  is 1) halogen,  
                   2) amino,  
                   3) carboxy,  
                   4)  $C_{1-4}$ alkyl,  
                   5)  $C_{1-4}$ alkoxy,

- 6) aryl,
- 7) aryl C<sub>1-4</sub>alkyl, or
- 8) aryloxy.

- 5 R<sup>d</sup> and R<sup>e</sup> are independently selected from hydrogen, C<sub>1-10</sub>alkyl, C<sub>2-10</sub>alkenyl, C<sub>2-10</sub>alkynyl, Cy and Cy C<sub>1-10</sub>alkyl, wherein alkyl, alkenyl, alkynyl and Cy is optionally substituted with one to four substituents independently selected from R<sup>c</sup>; or
- 10 R<sup>d</sup> and R<sup>e</sup> together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

R<sup>f</sup> and R<sup>g</sup> are independently selected from hydrogen, C<sub>1-10</sub>alkyl, Cy and Cy C<sub>1-10</sub>alkyl; or

- 15 R<sup>f</sup> and R<sup>g</sup> together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

- 20 R<sup>h</sup> is
- 1) hydrogen,
  - 2) C<sub>1-10</sub>alkyl,
  - 3) C<sub>2-10</sub>alkenyl,
  - 4) C<sub>2-10</sub>alkynyl,
  - 5) cyano,
  - 6) aryl,
  - 25 7) aryl C<sub>1-10</sub>alkyl,
  - 8) heteroaryl,
  - 9) heteroaryl C<sub>1-10</sub>alkyl, or
  - 10) -SO<sub>2</sub>R<sup>i</sup>;

- 30 wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from R<sup>a</sup>; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

- R<sup>i</sup> 1) C<sub>1-10</sub>alkyl,

- 2) C<sub>2</sub>-10alkenyl,
- 3) C<sub>2</sub>-10alkynyl, or
- 4) aryl;

wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted  
5 with one to four substituents independently selected from R<sup>c</sup>;

Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

m is an integer from 1 to 2;

10

n is an integer from 1 to 10;

- X is
- 1) -C(O)OR<sup>d</sup>,
  - 2) -P(O)(OR<sup>d</sup>)(OR<sup>e</sup>)
  - 15 3) -P(O)(R<sup>d</sup>)(OR<sup>e</sup>)
  - 4) -S(O)<sub>m</sub>OR<sup>d</sup>,
  - 5) -C(O)NR<sup>d</sup>R<sup>h</sup>, or
  - 6) -5-tetrazolyl;

- 20 Y is
- 1) -C(O)-,
  - 2) -O-C(O)-,
  - 3) -NR<sup>e</sup>-C(O)-,
  - 4) -S(O)<sub>2</sub>-,
  - 5) -P(O)(OR<sup>i</sup>)
  - 25 6) C(O)C(O).

2. A compound of Claim 1 wherein R<sup>1</sup> is C<sub>1</sub>-10alkyl, aryl, aryl-C<sub>1</sub>-10alkyl, heteroaryl or heteroaryl-C<sub>1</sub>-10alkyl, wherein alkyl, aryl and heteroaryl are optionally substituted as provided for in  
30 Claim 1.

3. A compound of Claim 2 wherein R<sup>1</sup> is phenyl optionally substituted with one to three groups selected from R<sup>b</sup>.

4. A compound of Claim 1 wherein R<sup>2</sup> is hydrogen, C<sub>1</sub>-10 alkyl, Cy or Cy-C<sub>1</sub>-10 alkyl; or R<sup>2</sup>, R<sup>3</sup> together with the atoms to which they are attached form a ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen, wherein said ring may be isolated or benzo-fused, and optionally substituted with one to four substituents independently selected from R<sup>b</sup>.

5. A compound of Claim 2 wherein R<sup>2</sup>, R<sup>3</sup> together with the atoms to which they are attached form a ring of 5 to 6 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen, wherein said ring may be isolated or benzo-fused, and optionally substituted with one to four substituents independently selected from R<sup>b</sup>.

6. A compound of Claim 1 wherein R<sup>4</sup> is H, C<sub>1</sub>-10alkyl, aryl, heteroaryl, aryl-C<sub>1</sub>-10alkyl or heteroaryl-C<sub>1</sub>-10alkyl. Preferably, R<sup>4</sup> is H or C<sub>1</sub>-10 alkyl.

7. A compound of Claim 6 wherein R<sup>4</sup> is H or C<sub>1</sub>-10alkyl.

8. A compound of Claim 1 wherein R<sup>6</sup> is Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>1</sub>-10alkyl wherein Ar<sup>1</sup> and Ar<sup>2</sup> are each aryl or heteroaryl each of which is optionally substituted with from 1 to 4 groups independently selected from R<sup>b</sup>.

9. A compound of Claim 8 wherein R<sup>6</sup> is Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>1</sub>-3alkyl wherein Ar<sup>1</sup> and Ar<sup>2</sup> are each aryl or heteroaryl each of which is optionally substituted with from 1 to 4 groups independently selected from R<sup>b</sup>.


10. A compound of Claim 1 wherein X is C(O)OR<sup>d</sup>.



11. A compound of Claim 1 wherein Y is C(O) or S(O)<sub>2</sub>.

12. A compound of Claim 11 wherein Y is S(O)<sub>2</sub>.

5 13. A compound of Claim 1 having the formula Ia:



wherein

- 10 R<sup>1</sup> is 1) C<sub>1</sub>-10alkyl,  
 2) Cy, or  
 3) Cy-C<sub>1</sub>-10alkyl,

wherein alkyl is optionally substituted with one to four substituents  
 independently selected from R<sup>a</sup>; and Cy is optionally substituted with  
 15 one to four substituents independently selected from R<sup>b</sup>;

- R<sup>4</sup> is 1) hydrogen, or  
 2) C<sub>1</sub>-3 alkyl;

Ar<sup>1</sup> and Ar<sup>2</sup> or independently aryl or heteroaryl each of which is  
 optionally substituted with one to four substituents independently  
 20 selected from R<sup>b</sup>; and

R<sup>b</sup> is as defined in Claim 1.

14. A compound of Claim 1 selected from the group  
 consisting of:

- 25 N-(3,4-dimethoxybenzenesulfonyl)-1,2,3,4-tetrahydro-isoquinoline-  
 3(S)-carbonyl-(L)-biphenylalanine;  
 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-biphenylalanine;  
 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4-  
 fluorophenyl)phenylalanine;  
 30 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-thienyl)-  
 phenylalanine;

- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-thienyl)-phenylalanine;
- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4'-trifluoromethyl-phenyl)-phenylalanine;
- 5 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methoxy-phenyl)-phenylalanine;
- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-formyl-phenyl)-phenylalanine;
- N-(3-fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-
- 10 thienyl)phenylalanine;
- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2',6'-difluorophenyl)phenylalanine;
- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-hydroxymethylphenyl)phenylalanine;
- 15 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4'-methylphenyl)phenylalanine;
- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-carboxyphenyl)phenylalanine;
- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-
- 20 methoxycarbonylphenyl)phenylalanine;
- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-formylphenyl)phenylalanine;
- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-aminophenyl)phenylalanine;
- 25 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylphenyl)phenylalanine;
- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-acetamidophenyl)phenylalanine;
- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-
- 30 fluorophenyl)phenylalanine;
- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-carboxyphenyl)phenylalanine;
- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-methoxycarbonylphenyl)phenylalanine;

- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2',4'-dichlorophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-formyl-3-thienyl)phenylalanine;  
5 N-(3,5-dichlorobenzenesulfonyl)-2(S)-methylprolyl-(L)-4-(4'-fluorophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-formylphenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-  
10 (hydroxymethyl)phenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(benzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-formylphenyl)phenylalanine;  
15 N-(benzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-methoxyphenyl)phenylalanine;  
N-(3-Fluorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(4'-fluoro-2'-  
20 methoxyphenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylthiophenyl)phenylalanine;  
25 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(5-methyl-1,3,4-oxadiazol-2-yl)-phenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-methyl-5-trifluoromethyl-benzoxazol-7-yl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-methyl-6-(5-  
30 trifluoromethyl-tetrazol-1-yl)-benzoxazol-4-yl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-methyl-5-(5-trifluoromethyl-tetrazol-1-yl)-benzoxazol-7-yl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3-pyridyl)phenylalanine;

- N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-pyridyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(5-pyrimidinyl)phenylalanine;  
5 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(3'-cyanophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2-methyl-benzoxazol-4-yl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(6-acetamido-2-methyl-benzoxazol-4-yl)phenylalanine;  
10 N-(benzenesulfonyl)-(L)-prolyl-(L)-4-(2-pyridyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-3(S)-methylprolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(tetrazol-5-yl)phenyl)phenylalanine;  
15 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(2-methyl-tetrazol-5-yl)phenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-(3-methyl-tetrazol-5-yl)phenyl)phenylalanine;  
20 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-aminocarbonylphenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylaminocarbonylphenyl)phenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
25 N-(benzenesulfonyl)-2(S)-methylprolyl-(L)-4-(2'-carboxyphenyl)phenylalanine;  
N-(3-bromobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
30 N-(benzenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-( $\alpha$ -toluenesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(phenylacetyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;;  
N-(3-pyridinesulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(2-thienylsulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;

- N-(benzylaminocarbonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-(3-phenylpropionyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-((5-methyl-3,4-thiadiazol-2-yl)sulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
5 N-((benzthiazol-2-yl)sulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
N-((1-methyl-imidazol-4-yl)sulfonyl)-(L)-prolyl-(L)-4-(2'-cyanophenyl)phenylalanine;  
10 N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylsulfinylphenyl)phenylalanine; and  
N-(3,5-dichlorobenzenesulfonyl)-(L)-prolyl-(L)-4-(2'-methylsulfonylphenyl)phenylalanine.

- 15           15. A method for inhibiting cell adhesion in a mammal which comprises administering to said mammal an effective amount of a compound of Claim 1.
- 20           16. A method for the treatment of diseases, disorders, conditions or symptoms mediated by cell adhesion in a mammal which comprises administering to said mammal an effective amount of a compound of Claim 1.
- 25           17. A method for the treatment of asthma, allergic rhinitis, multiple sclerosis, atherosclerosis or inflammation in a mammal which comprises administering to said mammal an effective amount of a compound of Claim 1.
- 30           18. A pharmaceutical composition which comprises a compound of Claim 1 and a pharmaceutically acceptable carrier thereof.
19. A method for inhibiting cell adhesion in a mammal which comprises administering to said mammal an effective amount of a compound of Claim 13.

20. A method for the treatment of diseases, disorders, conditions or symptoms mediated by cell adhesion in a mammal which comprises administering to said mammal an effective amount of a  
5 compound of Claim 13.

21. A method for the treatment of asthma, allergic rhinitis, multiple sclerosis, atherosclerosis or inflammation in a mammal which comprises administering to said mammal an effective amount of a  
10 compound of Claim 13.

22. A pharmaceutical composition which comprises a compound of Claim 13 and a pharmaceutically acceptable carrier thereof.  
15

**TITLE OF THE INVENTION**

**BIARYLALKANOIC ACID COMPOUNDS S AS CELL ADHESION  
INHIBITORS**

**5    ABSTRACT OF THE DISCLOSURE**

Biarylalkanoic acids of Formula I are antagonists of VLA-4, and as such are useful in the inhibition of prevention of cell adhesion and cell-adhesion mediated pathologies. These compounds may be formulated into pharmaceutical compositions and are suitable for use in  
10 the treatment of asthma, allergies, inflammation, multiple sclerosis, and other inflammatory and autoimmune disorders.

**THIS PAGE BLANK (USPTO)**

**THIS PAGE BLANK (USPTO)**